

2009

Does Calcium Interact With Titin's Immunoglobulin Domain in Cardiac Muscle?

DuVall, Michael

DuVall, M., Amrein, M., Gifford, J. & Herzog, W. "Does Calcium Interact With Titin's Immunoglobulin Domain in Cardiac Muscle?". 4th Annual Students' Union Undergraduate Research Symposium, November 18-19, 2009, University of Calgary, Calgary, AB.

<http://hdl.handle.net/1880/47634>

Downloaded from PRISM Repository, University of Calgary

Does Calcium Interact With Titin's Immunoglobulin Domain in Cardiac Muscle?



Michael DuVall, Matthias Amrein, Jessica Gifford, Walter Herzog
Human Performance Laboratory, The University of Calgary, Alberta, Canada



Introduction

- In North America, cardiac muscle diseases such as heart attacks and myopathies are on the rise.
- Contributing to knowledge in this area, we have focused on a critical muscle protein called titin (connectin) important in force production.
- Changes in the elastic function of cardiac titin regions accompany severe heart failure in humans [1].
- By adjusting the length of titin's extensible region, a muscle can vary its elastic properties [2] and thus passive force capability.
- Previous experiments with muscle fibers showed that titin-based tension is calcium responsive [3].

Purpose

- Identify the potential for passive force regulation of the immunoglobulin (Ig) domain within titin.
- Examine the effect of calcium on the chemical and structural properties of Ig domains.

Experimental Design

- E. Coli used to produce the 27th Ig domain from the I-band region of human cardiac titin. Proteins isolated using his-tagged affinity chromatography.
- Fluorescence spectroscopy was conducted on tryptophan π electrons using a Hitachi F-2000 wavelength scanning spectrofluorimeter at 295nm.




Figure 1: Schematic of titin isoform and arrangement of 8 tandem I27 domains. Ig (I): immunoglobulin; Fn (F): fibronectin; Single letters correspond to their respective amino acid or atomic abbreviations

- Atomic force microscopy (AFM) was performed using a JPK AFM mounted on a Zeiss inverted microscope.




Figure 2: Atomic force microscopy setup. Protein is attached to the substrate via covalent gold-sulfur linkages

Results




Figure 3: Silver stained SDS-PAGE of I27 domain separation with successively stronger imidazole washes (W)




Figure 4: Tryptophan fluorescence emission spectrum for titin I27 excited at 295nm with the addition of calcium (Ca^{2+}) and a calcium chelating agent (EDTA). Note the depression in intensity with increasing $[Ca^{2+}]$ and the partial reversal of this effect with EDTA




Figure 5: Force curve for I27 in relaxing solution indicating the force attained when the cantilever was extended toward (red) and retracted from (blue) the surface, fit to the Worm-Like Chain Model (green)




Figure 6: Three overlapping force curves for I27 in the presence of Ca^{2+} . Pulling speed: $1\mu m/sec$, $0.01M [Ca^{2+}]$. Note the gradual increase in force with Ca^{2+}

Discussion & Conclusion

- Fluorescence spectroscopy revealed a measurable degree of change in the microenvironment of I27 with the addition of calcium.
- Removing free calcium and repeating the fluorescence experiments lead to the almost complete restoration of fluorescence intensity, indicating reversibility.
- Tryptophan π electron delocalization suggests that calcium interacts and potentially binds to the Ig domain, which may have structural implications for titin's overall stiffness.
- Whether the alteration in the tryptophan environment translates into a change in the mechanical properties of I27, remains to be established.
- Using single molecule AFM, we have the means to evaluate differences in stiffness of Ig domains.
- The application of a mechanical force may trigger the exposure of new binding sites that were buried between domains or in the fold [4].
- Therefore, Ig domain unfolding within titin may modulate its resting length, elasticity and also its ligand binding properties [4].
- This entropic spring can then be used to better understand the changes in passive properties associated with specific cardiac failure.

References

- Neagoe C, et al. *Circulation*. 106, 1333-1341
- Linke WA, et al. *J Mol Biol* 261, 62-71
- Labeit D, et al. *Proc. Natl. Acad. Sci. U.S.A.* 2003. 100, 13716-13721
- Garcia TI, et al. *Proteins: Structure, Function and Bioinformatics* 75:3, 706-718

Acknowledgements

NSERC, USRP, PURE, Killam Foundation, Canada Research Chair Program, CIHR