Chapter 3

Reciprocal Regulation of Neuronal Calcium Channels by Synaptic Proteins

Norbert Weiss and Gerald W. Zamponi

Abstract Voltage-gated Ca^{2+} channels represent one of the main pathways for Ca^{2+} entry into nerve terminals where they play a critical role in the control of synaptic exocytosis. It is traditionally believed that the vesicle-docking/release machinery must be located in the vicinity of the calcium source in order to trigger fast, efficient and spatially delimited neurotransmitter release. This tight coupling is mostly achieved by a physical interaction of the presynaptic calcium channel with several actors of the synaptic vesicle release machinery. Conversely, the binding of synaptic proteins regulates calcium channel activity, providing for fine control of presynaptic Ca^{2+} entry. Here, we review the current state of knowledge of the molecular mechanisms by which synaptic proteins regulates presynaptic Ca^{2+} channel activity.

Keywords SNARE protein • Syntaxin • SNAP-25 • Synaptotagmin • Munc18 • Rim • Cysteine string proteins • Huntingtin

3.1 Introduction

Voltage-gated Ca^{2+} channels (VGCCs) are plasma membrane proteins that convert an electrical signal into intracellular Ca^{2+} elevations. To date, ten genes encoding the pore-forming subunits of mammalian VGCCs have been identified. Seven genes encode the high-voltage activated (HVA) channel subfamily (comprising L-type ($\text{Ca}_v1.1$ to $\text{Ca}_v1.4$), P/Q-type ($\text{Ca}_v2.1$), N-type ($\text{Ca}_v2.2$) and R-type ($\text{Ca}_v2.3$) channels) and three genes encode the low-voltage-activated (LVA) channel subfamily (composed exclusively of T-type ($\text{Ca}_v3.1$ to $\text{Ca}_v3.3$) (Ertel et al. 2000). In addition

N. Weiss (✉) • G.W. Zamponi

Department of Physiology and Pharmacology, Hotchkiss Brain Institute, University of Calgary, 3330 Hospital Dr. NW, Calgary, Canada T2N 4N1

e-mail: naweiss@ucalgary.ca; zamponi@ucalgary.ca

to the Ca_v pore-forming subunit, HVA channels contain auxiliary subunits: β (β_1 to β_4 , a 55 KDa cytosolic protein of the MAGUK (**m**embrane-**a**ssociated **g**uanylate **k**inase) family), $\alpha_2\delta$ ($\alpha_2\delta_1$ to $\alpha_2\delta_4$, a 170 KDa highly glycosylated extracellular protein with a single transmembrane domain), and in some cases γ (γ_1 to γ_8 , a 33 KDa transmembrane protein) (Takahashi et al. 1987). Among this wide diversity of native channels, $\text{Ca}_v2.1$ and $\text{Ca}_v2.2$ channels have been identified as the predominant Ca^{2+} channels involved in depolarization-evoked neurotransmitter release (Westenbroek et al. 1992, 1995, 1998; Olivera et al. 1994; Wheeler et al. 1994; Dunlap et al. 1995; Day et al. 1996; Timmermann et al. 2002). They support a transient Ca^{2+} microdomain of high concentration (10–50 μM) (Schneggenburger and Neher 2005) within the active zone of the synapse that is essential for the fusion of presynaptic vesicles with the plasma membrane (Llinas et al. 1992; Edwards 2007; Neher and Sakaba 2008; Weber et al. 2010). $\text{Ca}_v2.1$ channels support voltage-dependent exocytosis in the central nervous system, whereas $\text{Ca}_v2.2$ channels are critically involved at the peripheral synapses. In particular synapses, $\text{Ca}_v2.3$ channels are also expressed at sufficiently high density (Day et al. 1996; Hanson and Smith 2002) to support Ca^{2+} entry into presynaptic terminals (Breustedt et al. 2003; Dietrich et al. 2003) and contribute to some extent to the release of neurotransmitters (Wu et al. 1998; Gasparini et al. 2001; Kamp et al. 2005). Finally, although HVA channels support voltage-dependent exocytosis, release of neurotransmitters at rest (i.e. around the resting membrane potential of neurons) has been demonstrated in some neurons and relies on LVA $\text{Ca}_v3.2$ channels (Ivanov and Calabrese 2000; Pan et al. 2001; Egger et al. 2003; Carabelli et al. 2007; Weiss and Zamponi 2012).

In order to efficiently receive Ca^{2+} signals, the vesicle-docking/ release machinery must be located in the vicinity of the source of Ca^{2+} . This is particularly important considering the high Ca^{2+} buffering capability of neurons (Foehring et al. 2009) and the consequent limited diffusion of free Ca^{2+} . In mammalian synapses, this close localization relies on the direct interaction of the Ca^{2+} channels with several members of the vesicle release machinery which is essential for fast (within 200 μs after the arrival of the action potential) and spatially delimited neurotransmitter release (Sabatini and Regehr 1996; Wadel et al. 2007). Conversely, binding of synaptic proteins potently regulates channel activity, providing a reciprocal control of Ca^{2+} entry to fine tune synaptic strength. Interested readers may also refer to the work of Atlas et al., for a discussion of other possible role of biochemical coupling of VGCCs with synaptic proteins (Atlas et al. 2001; Lerner et al. 2006; Marom et al. 2007; Hagalili et al. 2008; Atlas 2010; Cohen-Kutner et al. 2010; Marom et al. 2010; Weiss 2010).

3.2 Basic Principles of Molecular Coupling Between Voltage-Gated Ca^{2+} Channels and SNARE Proteins

SNARE proteins (soluble NSF (*N*-ethylmaleimide-sensitive fusion protein) attachment protein receptor) comprising the Q-SNAREs syntaxin-1A/1B, SNAP-25 (synaptosomal-associated protein of 25 kDa) and R-SNARE synaptobrevin

(VAMP) (Fasshauer et al. 1998; Sutton et al. 1998) form the SNARE core complex that brings the vesicle and target membranes into close opposition, leading to fusion and exocytosis (Hanson et al. 1997; Otto et al. 1997). Not surprisingly, $\text{Ca}_v2.1$ and $\text{Ca}_v2.2$ channels are presynaptically colocalized with syntaxin-1A at nerve terminals (Cohen et al. 1991; Westenbroek et al. 1992, 1995) and have been biochemically isolated in complex with SNARE proteins (Bennett et al. 1992; Yoshida et al. 1992; Leveque et al. 1994). Molecular characterization of Ca^{2+} channels/SNARE interaction has identified a *synprint* (*synaptic protein interaction*) locus in $\text{Ca}_v2.1$ and $\text{Ca}_v2.2$ located within the intracellular loop between domains II and III of the channels (Sheng et al. 1994; Rettig et al. 1996). This motif binds syntaxin-1A and SNAP-25 (but not synaptobrevin). Further biochemical mapping of the *synprint* site has identified two distinct microdomains separated by a flexible linker that independently binds syntaxin-1A and SNAP-25 (Rettig et al. 1996; Yokoyama et al. 2005). The functional relevance of the interaction has been shown by disruption of the Ca^{2+} channel/SNAREs coupling using peptides derived from the *synprint* domain (or by direct deletion of the *synprint* site) that alters synaptic transmission (Mochida et al. 1996; Rettig et al. 1997; Harkins et al. 2004; Keith et al. 2007). However, although the *synprint* site is unambiguously of key importance for fast and efficient neurotransmitter release, there is evidence that some other channel isoforms such as T-type channels, although devoid of the consensus *synprint* site, functionally contribute to presynaptic Ca^{2+} elevations and neurotransmitter release, suggesting the existence of other molecular coupling determinants. Indeed, we recently demonstrated that syntaxin-1A and SNAP-25 biochemically interact with $\text{Ca}_v3.2$ T-type channels within the carboxy-terminal domain of the channel (Weiss et al. 2012; Fig. 3.1).

3.3 Functional Interaction of Voltage-Gated Ca^{2+} Channels with SNARE Proteins

SNARE proteins not only bring presynaptic vesicles close to the Ca^{2+} source but also potentially modulate channel gating to fine tune presynaptic Ca^{2+} entries and synaptic transmission (Fig. 3.2).

3.3.1 *Syntaxin-1A*

The notion that SNARE proteins modulate Ca^{2+} influx through VGCC arose from electrophysiological recordings in heterologous expression systems showing that coexpression of syntaxin-1A potentially modulates $\text{Ca}_v2.1$ and $\text{Ca}_v2.2$ gating by shifting the voltage-dependence of inactivation toward more negative membrane potentials (Bezprozvanny et al. 1995; Wiser et al. 1996; Zhong et al. 1999; Degtjar et al. 2000), thus silencing the channels and reducing presynaptic Ca^{2+} entry influx. This regulation was later confirmed in chick ciliary ganglion neurons and

Fig. 3.2 Functional interactions between presynaptic Ca^{2+} channels, synaptic proteins and second messengers. The SNARE syntaxin-1A and SNAP-25, as well as release of free G-protein $\beta\gamma$ dimer upon GPCR activation inhibit channel activity. In contrast, synaptotagmin-1, CSP and huntingtin prevent syntaxin-1A-dependent inhibition of the channel. Similarly, channel phosphorylation by PKC or CaM-KII prevents syntaxin-1A and $\text{G}\beta\gamma$ -dependent inhibition, while phosphorylation of SNAP-25 promotes its inhibitory effect. Rim-1 directly potentiates Ca^{2+} influx (Abbreviations: *Stx1A* syntaxin-1A, *Syt-1* synaptotagmin-1, *Rim-1* Rab-3 interacting molecule-1, *CSP* Cysteine String Proteins, *Htt* huntingtin, *PKC* protein kinase C, *CaM-KII* Ca^{2+} /calmodulin-dependent protein kinase II, *GPCR* G-protein coupled receptor. Arrows in red indicate an inhibitory regulation whereas arrows in green indicate a potentiation)

isolated mammalian nerve terminals (synaptosomes) upon application of botulinium neurotoxin C1 (BoNT/C1 which cleaves syntaxin-1A from its membrane anchoring domain). BoNT/C1 treatment shifted the voltage-dependence of inactivation of the channel toward depolarized potentials (Bergsman and Tsien 2000; Stanley 2003). Structure/function studies have identified the transmembrane domain of syntaxin-1A, in particularly the two cysteines (C271 and C272) (Trus et al. 2001), as well as a short stretch within the H3 helical cytoplasmic domain, as fundamental for channel modulation without a direct implication in biochemical interaction with the *synprint* domain (Bezprozvanny et al. 2000; Jarvis et al. 2002). More recently, it was also shown that besides binding the *synprint* site of $\text{Ca}_v2.2$ channels, the ten amino-terminal residues of syntaxin-1A might support inhibition of the channel (Davies et al. 2011). Although the exact molecular mechanism of syntaxin-1A mediated regulation of Ca^{2+} channels remains unclear, these results highlight the existence of two kinds of interaction of syntaxin-1A with the channel: (i) a biochemical interaction via the *synprint* domain and (ii) a functional interaction most likely involving additional yet unidentified channel determinants. Consistent with this idea, T-type Ca^{2+} channels that biochemically couple to syntaxin-1A

via different channel binding determinants than the consensus *synprint* domain, are subject to similar syntaxin-1A-dependent modulation as Cav2.1 and Cav2.2 channels. Similarly, for Cav2.3 channels, despite being devoid of the consensus *synprint* domain, various studies indicate that a similar syntaxin-1A-dependent regulation must occur (Bergsman and Tsien 2000; Wiser et al. 2002; Cohen and Atlas 2004). These observations suggest that gating modulation of VGCCs by syntaxin-1A likely involves modulatory channel determinants distinct from the anchoring domains. Dynamic intramolecular interactions between the intracellular loops of Cav2.1 and Cav2.2 channels have been reported and appear to be involved in channel gating behavior (Restituito et al. 2000; Raghib et al. 2001; Geib et al. 2002; Page et al. 2004, 2010; Sandoz et al. 2004; Agler et al. 2005; Bucci et al. 2011). Hence, channel remodeling upon syntaxin-1A binding might represent a possible consensus molecular mechanism by which syntaxin-1A modulates Cav2.x and Cav3.2 channels in a similar manner despite distinct coupling molecular determinants. Interestingly, a mutation (A454T) that segregates with familial hemiplegic migraine patients located within the intracellular linker between domains I and II of Cav2.1 channel alters both syntaxin-1A-dependent channel gating modulation and exocytosis (Serra et al. 2010), suggesting that the I-II channel loop could play an important role in mediating syntaxin-1A modulation. In addition, a conformational switch of syntaxin-1A has been reported, that depends of the molecular partners engaged in the macromolecular complex. Hence, syntaxin-1A presents a *closed* conformation in complex with munc18 (or in isolation) and switches to an *open* conformation when in complex with SNAP-25 or synaptobrevin-2 (Dulubova et al. 1999; Brunger 2001). Interestingly, whereas syntaxin-1A in its *closed* state potently modulates Cav2.2 and Cav3.2 channel activity, coexpression of an *open* syntaxin-1A (locked open by two point mutations (Dulubova et al. 1999)) no longer alters channel gating (Jarvis et al. 2002; Weiss et al. 2012). This conformational switch of syntaxin-1A appears to be of key physiological importance since it occurs during the vesicle release cycle (Dulubova et al. 1999; Fiebig et al. 1999; Richmond et al. 2001), suggesting that syntaxin-1A may dynamically and temporally controls presynaptic Ca²⁺ entry during the exocytosis process.

3.3.2 SNAP-25

Like syntaxin-1A, SNAP-25 non-competitively binds onto the *synprint* domain of Cav2.1 and Cav2.2 channels, as well as onto the carboxy-terminal domain of Cav3.2 channels, to produce a similar inhibitory channel gating modulation (Wiser et al. 1996; Zhong et al. 1999; Weiss et al. 2012). The functional modulation produced by SNAP-25 was recently indirectly confirmed in native systems where siRNA silencing of SNAP-25 in glutamatergic neurons produced an increase of Ca²⁺ currents carried by Cav2.1 channels due to a depolarizing shift of the voltage-dependence of inactivation (Condliffe et al. 2010; Condliffe and Matteoli 2011). Interestingly, this inhibitory regulation is no longer observed when SNAP-25 is

co-expressed with syntaxin-1A (Wiser et al. 1996; Zhong et al. 1999; Jarvis and Zamponi 2001a; Weiss et al. 2012), suggesting that association of SNAP-25 with syntaxin-1A during the vesicle release cycle relieves channel inhibition, allowing timely presynaptic Ca^{2+} elevation required for membrane fusion and exocytosis (Sudhof 2004). Consistent with a key physiological importance of SNAP-25, structural or expression alterations of the protein caused by genetic mutations have been associated with numerous neuropsychiatric and neurological disorders, likely because of mis-regulation of presynaptic Ca^{2+} channels (Corradini et al. 2009). Finally, it was reported that phosphorylation of SNAP-25 by protein kinase C is required for SNAP-25-dependent inhibition of VGCCs (Pozzi et al. 2008), suggesting that like syntaxin-1A, SNAP-25-dependent modulation of channel activity may involve molecular determinants other than the *synprint* site.

3.4 Modulation of Presynaptic Calcium Channels by Non SNARE Proteins

3.4.1 *Synaptotagmin-1*

Although part of the vesicular release complex, synaptotagmin-1 is not as essential as syntaxin-1A or SNAP-25 in the membrane fusion process per se (Tucker and Chapman 2002), but rather works as a Ca^{2+} sensor, forming the link between presynaptic Ca^{2+} elevation and vesicular fusion, essential for fast and synchronous neurotransmission release (DeBello et al. 1993; Geppert et al. 1994; Augustine 2001; Fernandez-Chacon et al. 2001; Nishiki and Augustine 2001; Tucker and Chapman 2002; Koh and Bellen 2003; Xu et al. 2007). Indeed, synaptotagmin-1 is characterized by an amino-terminal transmembrane region anchored in the vesicle, a variable linker, and two carboxy-terminal rich negatively charged domains (C2A and C2B), each capable of binding Ca^{2+} . Hence, Ca^{2+} binding onto the C2A domain contributes to the insertion of synaptotagmin-1 into the plasma membrane, bringing vesicles docked to the plasma membrane upon Ca^{2+} elevation (Fernandez-Chacon et al. 2001). In contrast, the C2B domain has been reported to biochemically interact with the *synprint* site of $\text{Ca}_v2.1$ and $\text{Ca}_v2.2$ channels (Sheng et al. 1997). Although binding of synaptotagmin-1 has no major effect on channel gating, it reduces syntaxin-1A-dependent inhibition of $\text{Ca}_v2.2$ channels, possibly by Ca^{2+} -dependent binding competition with syntaxin-1A (Sheng et al. 1996). Hence, syntaxin-1A preferentially interacts with the channel at rest (i.e. at low Ca^{2+} level) thus preventing channel activity, whereas presynaptic Ca^{2+} elevation favors its interaction with synaptotagmin-1 and Ca^{2+} entry through VGCCs that is required for the final fusion process of docked vesicles. Moreover, a Ca^{2+} -dependent synaptotagmin-1 interaction with the $\text{Ca}_v\beta_{4a}$ auxiliary-subunit of VGCCs has been reported (Vendel et al. 2006), providing another dynamic Ca^{2+} channel/vesicle interaction (Weiss 2006).

3.4.2 *Munc-18*

As for synaptotagmin-1, Munc18 belongs to the C2-domain containing protein family, and plays a fundamental role in the assembly/disassembly of the exocytosis machinery (Gulyas-Kovacs et al. 2007; Toonen and Verhage 2007). Genetic ablation of Munc-18 in mice leads to a complete loss of synaptic transmission (Verhage et al. 2000). Although a biochemical interaction of Munc-18 with Cav2.2 channel within the intracellular linker between domains II and III has been reported (Chan et al. 2007), its coexpression has no effect on channel gating (Gladysheva et al. 2004). Hence, Munc-18 appears to not be a direct modulator of Ca²⁺ channel activity but rather interferes with syntaxin-1A-dependent channel inhibition during the vesicle release cycle as previously mentioned (Dulubova et al. 1999; Brunger 2001). This occurs by stabilizing syntaxin-1A in a *closed* conformation (Jarvis et al. 2002), inhibiting Ca²⁺ channel activity and non-necessary presynaptic Ca²⁺ entry in the absence of docked vesicle.

3.4.3 *Rim-1*

Rim (Rab-3 interacting molecule) is also part of a family of vesicle-associated proteins whose members share C2 domains. By interacting with numerous components of the presynaptic active zone such as SNAP-25 or synaptotagmin-1 (Coppola et al. 2001), it forms a protein scaffold by participating in the docking and fusion of presynaptic vesicles (Wang et al. 2000; Betz et al. 2001; Coppola et al. 2001; Ohtsuka et al. 2002; Schoch et al. 2002; Kaeser et al. 2011). Essential for short- and long-term synaptic plasticity by affecting the readily releasable pool of vesicles (Castillo et al. 2002, 2002; Blundell et al. 2010; Deng et al. 2011; Han et al. 2011), Rim proteins are also essential for proper targeting of Ca²⁺ channels to presynaptic terminals (Han et al. 2011) and efficient neurotransmitter release (Schoch et al. 2006). Although biochemical studies using native synaptosome membrane preparations failed to demonstrate the existence of a Ca²⁺ channel/Rim complex (Wong and Stanley 2010), various studies report in vitro bindings of Rim with Ca²⁺ channel components. Indeed, direct interaction of Rim-1 with the *synprint* site of Cav2.2 channels has been shown (Coppola et al. 2001). Moreover, Rim Binding Proteins directly interact with Cav2.2 channels (and likely with Cav2.1 channels), possibly providing a molecular link between Ca²⁺ channels and Rim proteins (Hibino et al. 2002). Finally, biochemical interaction of Rim-1 with Cav β subunits has been reported, slowing Cav2.1, Cav2.2 and Cav2.3 channel inactivation when coexpressed in heterologous systems, thereby increasing Ca²⁺ influx during trains of action potentials (Kiyonaka et al. 2007), and a mutation in Rim-1 (R655H) associated with an autosomal dominant cone-rod dystrophy was found to alter Rim-1-dependent modulation of Cav2.1 channels gating (Miki et al. 2007) leading to a progressive loss of photoreceptors along with retinal degeneration (Barragan et al. 2005; Michaelides et al. 2005). Altogether, these results highlight the critical role of Rim-1 in the modulation of Ca²⁺ homeostasis at nerve terminals.

3.4.4 Cysteine String Proteins (CSP)

Cysteine String Proteins (CSP) are vesicle-associated protein with a key chaperone role at the synapse (Chamberlain and Burgoyne 2000). It was proposed that CSP may serve as a link between $\text{Ca}_v2.2$ channels and presynaptic vesicles (Mastrogia-como et al. 1994). Indeed, CSP interacts with the *synprint* motif of $\text{Ca}_v2.1$ (Leveque et al. 1998; Seagar et al. 1999; Magga et al. 2000) and $\text{Ca}_v2.2$ channels (Magga et al. 2000). Moreover CSP promotes presynaptic Ca^{2+} influx by recruiting dormant Ca^{2+} channels (Chen et al. 2002). Although the molecular mechanism by which CSP promotes channel activity remains unknown, considering that CSP interacts with syntaxin-1A (Nie et al. 1999; Wu et al. 1999), it is possible that binding of CSP onto syntaxin-1A prevents syntaxin-1A-dependent channel inhibition. Hence, like synaptotagmin-1, CSP may act as a molecular channel switch activity between undocked and docked vesicles release for timely control or presynaptic Ca^{2+} influx.

3.4.5 Huntingtin

Huntingtin (Htt) is well known for its implication in Huntington's disease (Ross and Tabrizi 2011) but the exact cellular function of the protein remains unclear. However, the observation that genetic ablation of Htt is lethal in mice highlights the fundamental importance of the protein (Nasir et al. 1995). Besides interacting with numerous proteins (to date at least 20 proteins involved in gene transcription, cellular transport or cell signaling has been shown to interact with Htt), Htt directly binds to the *synprint* domain of $\text{Ca}_v2.2$ channels (Swayne et al. 2005). However, as synaptotagmin-1, coexpression of Htt with $\text{Ca}_v2.2$ channels has no consequence on channel gating, but prevents syntaxin-1A-dependent regulation (Swayne et al. 2005), likely by displacing binding of syntaxin-1A from the channel (Swayne et al. 2006). Hence, Htt is not a direct channel modulator per se, but might represent an important actor of synaptic activity by influencing SNARE modulation. However, it remains unclear if Htt is permanently expressed at the synapse under normal condition, or if it is specifically targeted under particular physiopathological states.

3.5 Modulation of Calcium Channels by Other Signaling Pathways

3.5.1 G-Protein Coupled Receptors

Ca^{2+} entry into presynaptic terminals is also modulated by the activation of numerous G-protein-coupled receptors (GPCRs) and second messengers (Jarvis and Zamponi 2001b). Indeed, activation of specific GPCRs following the liberation of

neurotransmitters initiates a negative feedback regulation on presynaptic VGCCs, inhibiting presynaptic Ca^{2+} entry thus terminating synaptic transmission (Brown and Sihra 2008). This spatially delimited regulation (Forscher et al. 1986) relies on the direct binding of free G-protein $\beta\gamma$ dimers release upon GPCR activation (Herlitze et al. 1996; Ikeda 1996) to specific intracellular regions of the channel (De Waard et al. 1997, 2005; Zamponi et al. 1997; Tedford and Zamponi 2006). Interestingly, it was shown that cleavage of syntaxin-1A by the botulinium neurotoxin C1 in chick calyx synapses prevents G-protein-dependent inhibition of Ca^{2+} currents, suggesting the involvement of syntaxin-1A in presynaptic G-protein regulation of Ca^{2+} channels (Stanley and Mirotznik 1997; Silinsky 2005). In vitro studies have later revealed an interaction of syntaxin-1A with G-protein $\beta\gamma$ dimers, and although syntaxin-1A is not critical for G-protein regulation it potentiates the inhibition in a receptor-independent manner (Jarvis et al. 2000; Jarvis and Zamponi 2001a; Lu et al. 2001). Moreover, G-protein $\beta\gamma$ dimers not only interact with syntaxin-1A, but also with SNAP-25 to mediate presynaptic inhibition (Gerachshenko et al. 2005). Finally, besides to modulate Ca^{2+} channel activity, it was shown that G-protein $\beta\gamma$ dimers and synaptotagmin-1 compete for binding to the core SNARE complex in a Ca^{2+} -dependent manner such that at high Ca^{2+} concentration synaptotagmin-1 can displace $\text{G}\beta\gamma$ binding (Yoon et al. 2007). Hence, Ca^{2+} elevation in presynaptic terminals may prevent G-protein inhibition, likely by preventing binding of $\text{G}\beta\gamma$ with SNARE proteins (Yoon et al. 2007). To add further complexity, it was recently reported that Rim-1 promotes relief of G-protein inhibition of $\text{Ca}_v2.2$ channels by modulating channel inactivation (Weiss et al. 2011). Altogether, these results highlight the extreme interplay between GPCR-dependent regulation and the molecular actors of the exocytosis process to fine tune presynaptic Ca^{2+} entry.

3.5.2 Phosphorylation

In vitro studies have shown that the protein kinase C (PKC), as well as the Ca^{2+} -calmodulin-dependent kinase II (CaM-KII) are able to phosphorylate the *synprint* domain of $\text{Ca}_v2.2$ channels (Yokoyama et al. 1997, 2005), preventing binding of syntaxin-1A and SNAP-25 (Yokoyama et al. 1997) and thus preventing SNARE-dependent inhibition of the channel (Jarvis and Zamponi 2001a). However, uncoupling of SNARE proteins from the channel upon *synprint* phosphorylation, most likely represent a termination signal for synaptic exocytosis. It was reported that phosphorylation of the *synprint* site of $\text{Ca}_v2.1$ channels by the glycogen synthase kinase-3 (GSK-3) prevents SNARE interaction with the channel but also inhibits synaptic exocytosis possibly by interfering with Ca^{2+} channel/SNARE coupling (Zhu et al. 2010). Interestingly, phosphorylation of syntaxin-1A and SNAP-25 by PKC or CaM-KII does not alter interaction with the *synprint* site (Yokoyama et al. 1997). Hence, PKC- and CaM-KII-dependent phosphorylation of

the *synprint* site may serve as a biochemical switch for interaction/modulation of voltage-gated Ca^{2+} channels with SNARE protein complexes.

3.6 Concluding Remarks

Like most key cellular functions, control of neurotransmitter release by presynaptic Ca^{2+} channels is highly regulated. The various components of the exocytosis machinery, besides localizing the vesicles within the vicinity of the source of Ca^{2+} , provide a potent reciprocal control of presynaptic Ca^{2+} influx by modulating channel gating. This dynamic regulation appears to be fundamental to dynamically and temporally fine tune neurotransmitter release. Surprisingly, as highlighted in this chapter, the regulation of presynaptic Ca^{2+} channels appears extremely complex, with intricate interplay between different types of synaptic proteins and second messenger signaling pathways, but also highly redundant. This important redundancy in Ca^{2+} channel regulation by various presynaptic proteins might ensure a security control over a fundamental physiological function. Finally, although $\text{Cav}2.1/\text{Cav}2.2$ and $\text{Cav}3.2$ T-type channels use completely distinct channel molecular determinants to interact with the vesicular machinery, they are functionally regulated by syntaxin-1A and SNAP-25 in a strikingly similar manner. This may perhaps underscore the fundamental importance of localizing the exocytosis machinery near the source of Ca^{2+} , and providing tight control over Ca^{2+} entry.

Acknowledgements Norbert Weiss is supported by fellowships from Alberta Innovates—Health Solutions (AI-HS) and from Hotchkiss Brain Institute. Gerald W. Zamponi is funded by the Canadian Institutes of Health Research, is a Canada Research Chair and AI-HS Scientist.

References

- Agler HL, Evans J, Tay LH, Anderson MJ, Colecraft HM, Yue DT (2005) G protein-gated inhibitory module of N-type ($\text{Cav}2.2$) Ca^{2+} channels. *Neuron* 46:891–904
- Atlas D (2010) Signaling role of the voltage-gated calcium channel as the molecular on/off-switch of secretion. *Cell Signal* 22:1597–1603
- Atlas D, Wiser O, Trus M (2001) The voltage-gated Ca^{2+} channel is the Ca^{2+} sensor of fast neurotransmitter release. *Cell Mol Neurobiol* 21:717–731
- Augustine GJ (2001) How does calcium trigger neurotransmitter release? *Curr Opin Neurobiol* 11:320–326
- Barragan I, Marcos I, Borrego S, Antinolo G (2005) Molecular analysis of RIM1 in autosomal recessive Retinitis pigmentosa. *Ophthalmic Res* 37:89–93
- Bennett MK, Calakos N, Scheller RH (1992) Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* 257:255–259
- Bergsman JB, Tsien RW (2000) Syntaxin modulation of calcium channels in cortical synaptosomes as revealed by botulinum toxin C1. *J Neurosci* 20:4368–4378
- Betz A, Thakur P, Junge HJ, Ashery U, Rhee JS, Scheuss V et al (2001) Functional interaction of the active zone proteins Munc13-1 and RIM1 in synaptic vesicle priming. *Neuron* 30:183–196

- Bezprozvanny I, Scheller RH, Tsien RW (1995) Functional impact of syntaxin on gating of N-type and Q-type calcium channels. *Nature* 378:623–626
- Bezprozvanny I, Zhong P, Scheller RH, Tsien RW (2000) Molecular determinants of the functional interaction between syntaxin and N-type Ca^{2+} channel gating. *Proc Natl Acad Sci U S A* 97:13943–13948
- Blundell J, Kaeser PS, Sudhof TC, Powell CM (2010) RIM1alpha and interacting proteins involved in presynaptic plasticity mediate prepulse inhibition and additional behaviors linked to schizophrenia. *J Neurosci* 30:5326–5333
- Breustedt J, Vogt KE, Miller RJ, Nicoll RA, Schmitz D (2003) Alpha1E-containing Ca^{2+} channels are involved in synaptic plasticity. *Proc Natl Acad Sci U S A* 100:12450–12455
- Brown DA, Sihra TS (2008) Presynaptic signaling by heterotrimeric G-proteins. *Handb Exp Pharmacol* 184:207–260
- Brunner AT (2001) Structure of proteins involved in synaptic vesicle fusion in neurons. *Annu Rev Biophys Biomol Struct* 30:157–171
- Bucci G, Mochida S, Stephens GJ (2011) Inhibition of synaptic transmission and G protein modulation by synthetic $\text{Ca}_v2.2$ Ca^{2+} channel peptides. *J Physiol* 589:3085–3101
- Carabelli V, Marcantoni A, Comunanza V, de Luca A, Diaz J, Borges R et al (2007) Chronic hypoxia up-regulates alpha1H T-type channels and low-threshold catecholamine secretion in rat chromaffin cells. *J Physiol* 584:149–165
- Castillo PE, Schoch S, Schmitz F, Sudhof TC, Malenka RC (2002) RIM1alpha is required for presynaptic long-term potentiation. *Nature* 415:327–330
- Chamberlain LH, Burgoyne RD (2000) Cysteine-string protein: the chaperone at the synapse. *J Neurochem* 74:1781–1789
- Chan AW, Khanna R, Li Q, Stanley EF (2007) Munc18: a presynaptic transmitter release site N type ($\text{Ca}_v2.2$) calcium channel interacting protein. *Channels (Austin)* 1:11–20
- Chen S, Zheng X, Schulze KL, Morris T, Bellen H, Stanley EF (2002) Enhancement of presynaptic calcium current by cysteine string protein. *J Physiol* 538:383–389
- Cohen R, Atlas D (2004) R-type voltage-gated Ca^{2+} channel interacts with synaptic proteins and recruits synaptotagmin to the plasma membrane of *Xenopus* oocytes. *Neuroscience* 128:831–841
- Cohen MW, Jones OT, Angelides KJ (1991) Distribution of Ca^{2+} channels on frog motor nerve terminals revealed by fluorescent omega-conotoxin. *J Neurosci* 11:1032–1039
- Cohen-Kutner M, Nachmann D, Atlas D (2010) $\text{Ca}_v2.1$ (P/Q channel) interaction with synaptic proteins is essential for depolarization-evoked release. *Channels (Austin)* 4:266–277
- Condliffe SB, Matteoli M (2011) Inactivation kinetics of voltage-gated calcium channels in glutamatergic neurons are influenced by SNAP-25. *Channels (Austin)* 5:304–307
- Condliffe SB, Corradini I, Pozzi D, Verderio C, Matteoli M (2010) Endogenous SNAP-25 regulates native voltage-gated calcium channels in glutamatergic neurons. *J Biol Chem* 285:24968–24976
- Coppola T, Magnin-Luthi S, Perret-Menoud V, Gattesco S, Schiavo G, Regazzi R (2001) Direct interaction of the Rab3 effector RIM with Ca^{2+} channels, SNAP-25, and synaptotagmin. *J Biol Chem* 276:32756–32762
- Corradini I, Verderio C, Sala M, Wilson MC, Matteoli M (2009) SNAP-25 in neuropsychiatric disorders. *Ann N Y Acad Sci* 1152:93–99
- Davies JN, Zamponi GW (2008) Old proteins, developing roles: the regulation of calcium channels by synaptic proteins. *Channels (Austin)* 2:130–138
- Davies JN, Jarvis SE, Zamponi GW (2011) Bipartite syntaxin 1A interactions mediate $\text{Ca}_v2.2$ calcium channel regulation. *Biochem Biophys Res Commun* 411:562–568
- Day NC, Shaw PJ, McCormack AL, Craig PJ, Smith W, Beattie R et al (1996) Distribution of alpha 1A, alpha 1B and alpha 1E voltage-dependent calcium channel subunits in the human hippocampus and parahippocampal gyrus. *Neuroscience* 71:1013–1024
- De Waard M, Liu H, Walker D, Scott VE, Gurnett CA, Campbell KP (1997) Direct binding of G-protein betagamma complex to voltage-dependent calcium channels. *Nature* 385:446–450

- De Waard M, Hering J, Weiss N, Feltz A (2005) How do G proteins directly control neuronal Ca^{2+} channel function? *Trends Pharmacol Sci* 26:427–436
- DeBello WM, Betz H, Augustine GJ (1993) Synaptotagmin and neurotransmitter release. *Cell* 74:947–950
- Degtiar VE, Scheller RH, Tsien RW (2000) Syntaxin modulation of slow inactivation of N-type calcium channels. *J Neurosci* 20:4355–4367
- Deng L, Kaeser PS, Xu W, Sudhof TC (2011) RIM proteins activate vesicle priming by reversing autoinhibitory homodimerization of Munc13. *Neuron* 69:317–331
- Dietrich D, Kirschstein T, Kukley M, Pereverzev A, von der Brölie C, Schneider T et al (2003) Functional specialization of presynaptic $\text{Ca}_v2.3$ Ca^{2+} channels. *Neuron* 39:483–496
- Dulubova I, Sugita S, Hill S, Hosaka M, Fernandez I, Sudhof TC et al (1999) A conformational switch in syntaxin during exocytosis: role of munc18. *Embo J* 18:4372–4382
- Dunlap K, Luebke JI, Turner TJ (1995) Exocytotic Ca^{2+} channels in mammalian central neurons. *Trends Neurosci* 18:89–98
- Edwards RH (2007) The neurotransmitter cycle and quantal size. *Neuron* 55:835–858
- Enger V, Svoboda K, Mainen ZF (2003) Mechanisms of lateral inhibition in the olfactory bulb: efficiency and modulation of spike-evoked calcium influx into granule cells. *J Neurosci* 23:7551–7558
- Ertel EA, Campbell KP, Harpold MM, Hofmann F, Mori Y, Perez-Reyes E et al (2000) Nomenclature of voltage-gated calcium channels. *Neuron* 25:533–535
- Fasshauer D, Sutton RB, Brunger AT, Jahn R (1998) Conserved structural features of the synaptic fusion complex: SNARE proteins reclassified as Q- and R-SNAREs. *Proc Natl Acad Sci U S A* 95:15781–15786
- Fernandez-Chacon R, Konigstorfer A, Gerber SH, Garcia J, Matos MF, Stevens CF et al (2001) Synaptotagmin I functions as a calcium regulator of release probability. *Nature* 410:41–49
- Fiebig KM, Rice LM, Pollock E, Brunger AT (1999) Folding intermediates of SNARE complex assembly. *Nat Struct Biol* 6:117–123
- Foehring RC, Zhang XF, Lee JC, Callaway JC (2009) Endogenous calcium buffering capacity of substantia nigral dopamine neurons. *J Neurophysiol* 102:2326–2333
- Forscher P, Oxford GS, Schulz D (1986) Noradrenaline modulates calcium channels in avian dorsal root ganglion cells through tight receptor-channel coupling. *J Physiol* 379:131–144
- Gasparini S, Kasyanov AM, Pietrobon D, Voronin LL, Cherubini E (2001) Presynaptic R-type calcium channels contribute to fast excitatory synaptic transmission in the rat hippocampus. *J Neurosci* 21:8715–8721
- Geib S, Sandoz G, Cornet V, Mabrouk K, Fund-Saunier O, Bichet D et al (2002) The interaction between the I-II loop and the III-IV loop of $\text{Ca}_v2.1$ contributes to voltage-dependent inactivation in a β -dependent manner. *J Biol Chem* 277:10003–10013
- Geppert M, Goda Y, Hammer RE, Li C, Rosahl TW, Stevens CF et al (1994) Synaptotagmin I: a major Ca^{2+} sensor for transmitter release at a central synapse. *Cell* 79:717–727
- Gerachshenko T, Blackmer T, Yoon EJ, Bartleson C, Hamm HE, Alford S (2005) Gbetagamma acts at the C terminus of SNAP-25 to mediate presynaptic inhibition. *Nat Neurosci* 8:597–605
- Gladysheva SE, Ho CS, Lee YY, Stuenkel EL (2004) Regulation of syntaxin1A-munc18 complex for SNARE pairing in HEK293 cells. *J Physiol* 558:857–871
- Gulyas-Kovacs A, de Wit H, Milosevic I, Kochubey O, Toonen R, Klingauf J et al (2007) Munc18-1: sequential interactions with the fusion machinery stimulate vesicle docking and priming. *J Neurosci* 27:8676–8686
- Hagalili Y, Bachnoff N, Atlas D (2008) The voltage-gated Ca^{2+} channel is the Ca^{2+} sensor protein of secretion. *Biochemistry* 47:13822–13830
- Han Y, Kaeser PS, Sudhof TC, Schneggenburger R (2011) RIM determines Ca^{2+} channel density and vesicle docking at the presynaptic active zone. *Neuron* 69:304–316
- Hanson JE, Smith Y (2002) Subcellular distribution of high-voltage-activated calcium channel subtypes in rat globus pallidus neurons. *J Comp Neurol* 442:89–98
- Hanson PI, Heuser JE, Jahn R (1997) Neurotransmitter release—four years of SNARE complexes. *Curr Opin Neurobiol* 7:310–315

- Harkins AB, Cahill AL, Powers JF, Tischler AS, Fox AP (2004) Deletion of the synaptic protein interaction site of the N-type ($\text{Ca}_v2.2$) calcium channel inhibits secretion in mouse pheochromocytoma cells. *Proc Natl Acad Sci U S A* 101:15219–15224
- Herlitze S, Garcia DE, Mackie K, Hille B, Scheuer T, Catterall WA (1996) Modulation of Ca^{2+} channels by G-protein beta gamma subunits. *Nature* 380:258–262
- Hibino H, Pironkova R, Onwumere O, Vologodskaya M, Hudspeth AJ, Lesage F (2002) RIM binding proteins (RBPs) couple Rab3-interacting molecules (RIMs) to voltage-gated Ca^{2+} channels. *Neuron* 34:411–423
- Ikeda SR (1996) Voltage-dependent modulation of N-type calcium channels by G-protein beta gamma subunits. *Nature* 380:255–258
- Ivanov AI, Calabrese RL (2000) Intracellular Ca^{2+} dynamics during spontaneous and evoked activity of leech heart interneurons: low-threshold Ca currents and graded synaptic transmission. *J Neurosci* 20:4930–4943
- Jarvis SE, Zamponi GW (2001a) Distinct molecular determinants govern syntaxin 1A-mediated inactivation and G-protein inhibition of N-type calcium channels. *J Neurosci* 21:2939–2948
- Jarvis SE, Zamponi GW (2001b) Interactions between presynaptic Ca^{2+} channels, cytoplasmic messengers and proteins of the synaptic vesicle release complex. *Trends Pharmacol Sci* 22: 519–525
- Jarvis SE, Magga JM, Beedle AM, Braun JE, Zamponi GW (2000) G protein modulation of N-type calcium channels is facilitated by physical interactions between syntaxin 1A and Gbetagamma. *J Biol Chem* 275:6388–6394
- Jarvis SE, Barr W, Feng ZP, Hamid J, Zamponi GW (2002) Molecular determinants of syntaxin 1 modulation of N-type calcium channels. *J Biol Chem* 277:44399–44407
- Kaesler PS, Deng L, Wang Y, Dulubova I, Liu X, Rizo J et al (2011) RIM proteins tether Ca^{2+} channels to presynaptic active zones via a direct PDZ-domain interaction. *Cell* 144:282–295
- Kamp MA, Krieger A, Henry M, Hescheler J, Weiergraber M, Schneider T (2005) Presynaptic ' $\text{Ca}_v2.3$ -containing' E-type Ca channels share dual roles during neurotransmitter release. *Eur J Neurosci* 21:1617–1625
- Keith RK, Poage RE, Yokoyama CT, Catterall WA, Meriney SD (2007) Bidirectional modulation of transmitter release by calcium channel/syntaxin interactions in vivo. *J Neurosci* 27:265–269
- Kiyonaka S, Wakamori M, Miki T, Uriu Y, Nonaka M, Bito H et al (2007) RIM1 confers sustained activity and neurotransmitter vesicle anchoring to presynaptic Ca^{2+} channels. *Nat Neurosci* 10:691–701
- Koh TW, Bellen HJ (2003) Synaptotagmin I, a Ca^{2+} sensor for neurotransmitter release. *Trends Neurosci* 26:413–422
- Lerner I, Trus M, Cohen R, Yizhar O, Nussinovitch I, Atlas D (2006) Ion interaction at the pore of Lc-type Ca^{2+} channel is sufficient to mediate depolarization-induced exocytosis. *J Neurochem* 97:116–127
- Leveque C, el Far O, Martin-Moutot N, Sato K, Kato R, Takahashi M et al (1994) Purification of the N-type calcium channel associated with syntaxin and synaptotagmin. A complex implicated in synaptic vesicle exocytosis. *J Biol Chem* 269:6306–6312
- Leveque C, Pupier S, Marqueze B, Geslin L, Kataoka M, Takahashi M et al (1998) Interaction of cysteine string proteins with the alpha1A subunit of the P/Q-type calcium channel. *J Biol Chem* 273:13488–13492
- Llinas R, Sugimori M, Silver RB (1992) Microdomains of high calcium concentration in a presynaptic terminal. *Science* 256:677–679
- Lu Q, AtKisson MS, Jarvis SE, Feng ZP, Zamponi GW, Dunlap K (2001) Syntaxin 1A supports voltage-dependent inhibition of alpha1B Ca^{2+} channels by Gbetagamma in chick sensory neurons. *J Neurosci* 21:2949–2957
- Magga JM, Jarvis SE, Arnot MI, Zamponi GW, Braun JE (2000) Cysteine string protein regulates G protein modulation of N-type calcium channels. *Neuron* 28:195–204
- Marom M, Sebag A, Atlas D (2007) Cations residing at the selectivity filter of the voltage-gated Ca^{2+} -channel modify fusion-pore kinetics. *Channels (Austin)* 1:377–386

- Marom M, Hagalili Y, Sebag A, Tzvier L, Atlas D (2010) Conformational changes induced in voltage-gated calcium channel $\text{Ca}_v1.2$ by BayK 8644 or FPL64176 modify the kinetics of secretion independently of Ca^{2+} influx. *J Biol Chem* 285:6996–7005
- Mastrogiacomo A, Parsons SM, Zampighi GA, Jenden DJ, Umbach JA, Gundersen CB (1994) Cysteine string proteins: a potential link between synaptic vesicles and presynaptic Ca^{2+} channels. *Science* 263:981–982
- Michaelides M, Holder GE, Hunt DM, Fitzke FW, Bird AC, Moore AT (2005) A detailed study of the phenotype of an autosomal dominant cone-rod dystrophy (CORD7) associated with mutation in the gene for RIM1. *Br J Ophthalmol* 89:198–206
- Miki T, Kiyonaka S, Uriu Y, De Waard M, Wakamori M, Beedle AM et al (2007) Mutation associated with an autosomal dominant cone-rod dystrophy CORD7 modifies RIM1-mediated modulation of voltage-dependent Ca^{2+} channels. *Channels (Austin)* 1:144–147
- Mochida S, Sheng ZH, Baker C, Kobayashi H, Catterall WA (1996) Inhibition of neurotransmission by peptides containing the synaptic protein interaction site of N-type Ca^{2+} channels. *Neuron* 17:781–788
- Nasir J, Floresco SB, O’Kusky JR, Diewert VM, Richman JM, Zeisler J et al (1995) Targeted disruption of the Huntington’s disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes. *Cell* 81:811–823
- Neher E, Sakaba T (2008) Multiple roles of calcium ions in the regulation of neurotransmitter release. *Neuron* 59:861–872
- Nie Z, Ranjan R, Wenniger JJ, Hong SN, Bronk P, Zinsmaier KE (1999) Overexpression of cysteine-string proteins in *Drosophila* reveals interactions with syntaxin. *J Neurosci* 19:10270–10279
- Nishiki TI, Augustine GJ (2001) Calcium-dependent neurotransmitter release: synaptotagmin to the rescue. *J Comp Neurol* 436:1–3
- Ohtsuka T, Takao-Rikitsu E, Inoue E, Inoue M, Takeuchi M, Matsubara K et al (2002) Cast: a novel protein of the cytomatrix at the active zone of synapses that forms a ternary complex with RIM1 and munc13-1. *J Cell Biol* 158:577–590
- Olivera BM, Miljanich GP, Ramachandran J, Adams ME (1994) Calcium channel diversity and neurotransmitter release: the omega-conotoxins and omega-agatoxins. *Annu Rev Biochem* 63:823–867
- Otto H, Hanson PI, Jahn R (1997) Assembly and disassembly of a ternary complex of synaptobrevin, syntaxin, and SNAP-25 in the membrane of synaptic vesicles. *Proc Natl Acad Sci U S A* 94:6197–6201
- Page KM, Hebllich F, Davies A, Butcher AJ, Leroy J, Bertaso F et al (2004) Dominant-negative calcium channel suppression by truncated constructs involves a kinase implicated in the unfolded protein response. *J Neurosci* 24:5400–5409
- Page KM, Hebllich F, Margas W, Pratt WS, Nieto-Rostro M, Chaggar K et al (2010) N terminus is key to the dominant negative suppression of Ca_v2 calcium channels: implications for episodic ataxia type 2. *J Biol Chem* 285:835–844
- Pan ZH, Hu HJ, Perring P, Andrade R (2001) T-type Ca^{2+} channels mediate neurotransmitter release in retinal bipolar cells. *Neuron* 32:89–98
- Pozzi D, Condliffe S, Bozzi Y, Chikhladze M, Grumelli C, Proux-Gillardeaux V et al (2008) Activity-dependent phosphorylation of Ser187 is required for SNAP-25-negative modulation of neuronal voltage-gated calcium channels. *Proc Natl Acad Sci U S A* 105:323–328
- Raghib A, Bertaso F, Davies A, Page KM, Meir A, Bogdanov Y et al (2001) Dominant-negative synthesis suppression of voltage-gated calcium channel $\text{Ca}_v2.2$ induced by truncated constructs. *J Neurosci* 21:8495–8504
- Restituto S, Cens T, Barrere C, Geib S, Galas S, De Waard M et al (2000) The $\beta 2a$ subunit is a molecular groom for the Ca^{2+} channel inactivation gate. *J Neurosci* 20:9046–9052
- Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, Catterall WA (1996) Isoform-specific interaction of the $\alpha 1A$ subunits of brain Ca^{2+} channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci U S A* 93:7363–7368

- Rettig J, Heinemann C, Ashery U, Sheng ZH, Yokoyama CT, Catterall WA et al (1997) Alteration of Ca^{2+} dependence of neurotransmitter release by disruption of Ca^{2+} channel/syntaxin interaction. *J Neurosci* 17:6647–6656
- Richmond JE, Weimer RM, Jorgensen EM (2001) An open form of syntaxin bypasses the requirement for UNC-13 in vesicle priming. *Nature* 412:338–341
- Ross CA, Tabrizi SJ (2011) Huntington's disease: from molecular pathogenesis to clinical treatment. *Lancet Neurol* 10:83–98
- Sabatini BL, Regehr WG (1996) Timing of neurotransmission at fast synapses in the mammalian brain. *Nature* 384:170–172
- Sandoz G, Lopez-Gonzalez I, Stambouliau S, Weiss N, Arnoult C, De Waard M (2004) Repositioning of charged I-II loop amino acid residues within the electric field by beta subunit as a novel working hypothesis for the control of fast P/Q calcium channel inactivation. *Eur J Neurosci* 19:1759–1772
- Schneggenburger R, Neher E (2005) Presynaptic calcium and control of vesicle fusion. *Curr Opin Neurobiol* 15:266–274
- Schoch S, Castillo PE, Jo T, Mukherjee K, Geppert M, Wang Y et al (2002) RIM1alpha forms a protein scaffold for regulating neurotransmitter release at the active zone. *Nature* 415:321–326
- Schoch S, Mittelstaedt T, Kaeser PS, Padgett D, Feldmann N, Chevaleyre V et al (2006) Redundant functions of RIM1alpha and RIM2alpha in Ca^{2+} -triggered neurotransmitter release. *Embo J* 25:5852–5863
- Seagar M, Leveque C, Charvin N, Marqueze B, Martin-Moutot N, Boudier JA et al (1999) Interactions between proteins implicated in exocytosis and voltage-gated calcium channels. *Philos Trans R Soc Lond B Biol Sci* 354:289–297
- Serra SA, Cuenca-Leon E, Llobet A, Rubio-Moscardo F, Plata C, Carreno O et al (2010) A mutation in the first intracellular loop of CACNA1A prevents P/Q channel modulation by SNARE proteins and lowers exocytosis. *Proc Natl Acad Sci U S A* 107:1672–1677
- Sheng ZH, Rettig J, Takahashi M, Catterall WA (1994) Identification of a syntaxin-binding site on N-type calcium channels. *Neuron* 13:1303–1313
- Sheng ZH, Rettig J, Cook T, Catterall WA (1996) Calcium-dependent interaction of N-type calcium channels with the synaptic core complex. *Nature* 379:451–454
- Sheng ZH, Yokoyama CT, Catterall WA (1997) Interaction of the synprint site of N-type Ca^{2+} channels with the C2B domain of synaptotagmin I. *Proc Natl Acad Sci U S A* 94:5405–5410
- Silinsky EM (2005) Modulation of calcium currents is eliminated after cleavage of a strategic component of the mammalian secretory apparatus. *J Physiol* 566:681–688
- Stanley EF (2003) Syntaxin I modulation of presynaptic calcium channel inactivation revealed by botulinum toxin C1. *Eur J Neurosci* 17:1303–1305
- Stanley EF, Mirotznik RR (1997) Cleavage of syntaxin prevents G-protein regulation of presynaptic calcium channels. *Nature* 385:340–343
- Sudhof TC (2004) The synaptic vesicle cycle. *Annu Rev Neurosci* 27:509–547
- Sutton RB, Fasshauer D, Jahn R, Brunger AT (1998) Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* 395:347–353
- Swayne LA, Chen L, Hameed S, Barr W, Charlesworth E, Colicos MA et al (2005) Crosstalk between huntingtin and syntaxin 1A regulates N-type calcium channels. *Mol Cell Neurosci* 30:339–351
- Swayne LA, Beck KE, Braun JE (2006) The cysteine string protein multimeric complex. *Biochem Biophys Res Commun* 348:83–91
- Takahashi M, Seagar MJ, Jones JF, Reber BF, Catterall WA (1987) Subunit structure of dihydropyridine-sensitive calcium channels from skeletal muscle. *Proc Natl Acad Sci U S A* 84:5478–5482
- Tedford HW, Zamponi GW (2006) Direct G protein modulation of Ca_v2 calcium channels. *Pharmacol Rev* 58:837–862
- Timmermann DB, Westenbroek RE, Schousboe A, Catterall WA (2002) Distribution of high-voltage-activated calcium channels in cultured gamma-aminobutyric acidergic neurons from mouse cerebral cortex. *J Neurosci Res* 67:48–61

- Toonen RF, Verhage M (2007) Munc18-1 in secretion: lonely Munc joins SNARE team and takes control. *Trends Neurosci* 30:564–572
- Trus M, Wiser O, Goodnough MC, Atlas D (2001) The transmembrane domain of syntaxin 1A negatively regulates voltage-sensitive Ca^{2+} channels. *Neuroscience* 104:599–607
- Tucker WC, Chapman ER (2002) Role of synaptotagmin in Ca^{2+} -triggered exocytosis. *Biochem J* 366:1–13
- Vendel AC, Terry MD, Striegel AR, Iverson NM, Leuranguer V, Rithner CD et al (2006) Alternative splicing of the voltage-gated Ca^{2+} channel beta4 subunit creates a uniquely folded N-terminal protein binding domain with cell-specific expression in the cerebellar cortex. *J Neurosci* 26:2635–2644
- Verhage M, Maia AS, Plomp JJ, Brussaard AB, Heeroma JH, Vermeer H et al (2000) Synaptic assembly of the brain in the absence of neurotransmitter secretion. *Science* 287:864–869
- Wadel K, Neher E, Sakaba T (2007) The coupling between synaptic vesicles and Ca^{2+} channels determines fast neurotransmitter release. *Neuron* 53:563–575
- Wang Y, Sugita S, Sudhof TC (2000) The RIM/NIM family of neuronal C2 domain proteins. Interactions with Rab3 and a new class of Src homology 3 domain proteins. *J Biol Chem* 275:20033–20044
- Weber AM, Wong FK, Tufford AR, Schlichter LC, Matveev V, Stanley EF (2010) N-type Ca^{2+} channels carry the largest current: implications for nanodomains and transmitter release. *Nat Neurosci* 13:1348–1350
- Weiss N (2006) The calcium channel beta4a subunit: a scaffolding protein between voltage-gated calcium channel and presynaptic vesicle-release machinery? *J Neurosci* 26:6117–6118
- Weiss N (2010) Control of depolarization-evoked presynaptic neurotransmitter release by $\text{Ca}_v2.1$ calcium channel: old story, new insights. *Channels (Austin)* 4:431–433
- Weiss N, Zamponi GW (2012) Control of low-threshold exocytosis by T-type calcium channels. *Biochim Biophys Acta* 85:131–136
- Weiss N, Sandoval A, Kyonaka S, Felix R, Mori Y, De Waard M (2011) Rim1 modulates direct G-protein regulation of $\text{Ca}_v2.2$ channels. *Pflugers Arch* 461:447–459
- Weiss N, Hameed S, Fernandez-Fernandez JM, Fablet K, Karmazinova M, Poillot C et al (2012) A $\text{Ca}_v3.2$ /syntaxin-1A signaling complex controls T-type channel activity and low-threshold exocytosis. *J Biol Chem* 287:2810–2818
- Westenbroek RE, Hell JW, Warner C, Dubel SJ, Snutch TP, Catterall WA (1992) Biochemical properties and subcellular distribution of an N-type calcium channel alpha 1 subunit. *Neuron* 9:1099–1115
- Westenbroek RE, Sakurai T, Elliott EM, Hell JW, Starr TV, Snutch TP et al (1995) Immunohistochemical identification and subcellular distribution of the alpha 1A subunits of brain calcium channels. *J Neurosci* 15:6403–6418
- Westenbroek RE, Hoskins L, Catterall WA (1998) Localization of Ca^{2+} channel subtypes on rat spinal motor neurons, interneurons, and nerve terminals. *J Neurosci* 18:6319–6330
- Wheeler DB, Randall A, Tsien RW (1994) Roles of N-type and Q-type Ca^{2+} channels in supporting hippocampal synaptic transmission. *Science* 264:107–111
- Wiser O, Bennett MK, Atlas D (1996) Functional interaction of syntaxin and SNAP-25 with voltage-sensitive L- and N-type Ca^{2+} channels. *Embo J* 15:4100–4110
- Wiser O, Cohen R, Atlas D (2002) Ionic dependence of Ca^{2+} channel modulation by syntaxin 1A. *Proc Natl Acad Sci U S A* 99:3968–3973
- Wong FK, Stanley EF (2010) Rab3a interacting molecule (RIM) and the tethering of pre-synaptic transmitter release site-associated $\text{Ca}_v2.2$ calcium channels. *J Neurochem* 112:463–473
- Wu LG, Borst JG, Sakmann B (1998) R-type Ca^{2+} currents evoke transmitter release at a rat central synapse. *Proc Natl Acad Sci U S A* 95:4720–4725
- Wu MN, Fergestad T, Lloyd TE, He Y, Broadie K, Bellen HJ (1999) Syntaxin 1A interacts with multiple exocytic proteins to regulate neurotransmitter release in vivo. *Neuron* 23:593–605
- Xu J, Mashimo T, Sudhof TC (2007) Synaptotagmin-1, -2, and -9: Ca^{2+} sensors for fast release that specify distinct presynaptic properties in subsets of neurons. *Neuron* 54:567–581

- Yokoyama CT, Sheng ZH, Catterall WA (1997) Phosphorylation of the synaptic protein interaction site on N-type calcium channels inhibits interactions with SNARE proteins. *J Neurosci* 17:6929–6938
- Yokoyama CT, Myers SJ, Fu J, Mockus SM, Scheuer T, Catterall WA (2005) Mechanism of SNARE protein binding and regulation of Cav2 channels by phosphorylation of the synaptic protein interaction site. *Mol Cell Neurosci* 28:1–17
- Yoon EJ, Gerachshenko T, Spiegelberg BD, Alford S, Hamm HE (2007) Gbetagamma interferes with Ca^{2+} -dependent binding of synaptotagmin to the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex. *Mol Pharmacol* 72:1210–1219
- Yoshida A, Oho C, Omori A, Kuwahara R, Ito T, Takahashi M (1992) HPC-1 is associated with synaptotagmin and omega-conotoxin receptor. *J Biol Chem* 267:24925–24928
- Zamponi GW, Bourinet E, Nelson D, Nargeot J, Snutch TP (1997) Crosstalk between G proteins and protein kinase C mediated by the calcium channel $\alpha 1$ subunit. *Nature* 385:442–446
- Zhong H, Yokoyama CT, Scheuer T, Catterall WA (1999) Reciprocal regulation of P/Q-type Ca^{2+} channels by SNAP-25, syntaxin and synaptotagmin. *Nat Neurosci* 2:939–941
- Zhu LQ, Liu D, Hu J, Cheng J, Wang SH, Wang Q et al (2010) GSK-3 beta inhibits presynaptic vesicle exocytosis by phosphorylating P/Q-type calcium channel and interrupting SNARE complex formation. *J Neurosci* 30:3624–3633