UNIVERSITY OF CALGARY

Stratigraphy, Structure and Tectonic History of the Pink Mountain Anticline,
Trutch (94G) and Halfway River (94B) Map Areas, Northeastern British Columbia

by

Steven Jeffrey Hinds

A THESIS SUBMITTED TO THE FACULTY OF GRADUATE
STUDIES IN PARTIAL FULFILMENT OF THE REQUIREMENTS
FOR THE DEGREE OF MASTER OF SCIENCE

DEPARTMENT OF GEOLOGY AND GEOPHYSICS

CALGARY, ALBERTA

JUNE, 2002

© Steven Jeffrey Hinds 2002

UNIVERSITY OF
CALGARY

The author of this thesis has granted the University of Calgary a non-exclusive license to reproduce and distribute copies of this thesis to users of the University of Calgary Archives.

Copyright remains with the author.

Theses and dissertations available in the University of Calgary Institutional Repository are solely for the purpose of private study and research. They may not be copied or reproduced, except as permitted by copyright laws, without written authority of the copyright owner. Any commercial use or re-publication is strictly prohibited.

The original Partial Copyright License attesting to these terms and signed by the author of this thesis may be found in the original print version of the thesis, held by the University of Calgary Archives.

Please contact the University of Calgary Archives for further information:

E-mail: uarc@ucalgary.ca

Telephone: (403) 220-7271

Website: <http://archives.ucalgary.ca>

ABSTRACT

Pink Mountain Anticline stands out in front of the Foothills of northeastern British Columbia (57°N, 123°W). Geologic mapping and prestack depth-migrated seismic sections show that it is localized above and west of a northwest-trending subsurface normal fault. Along with isopach maps they demonstrate episodic normal movement during deposition of the Carboniferous Stoddart Group, Triassic Montney Formation and possibly the Jurassic-Cretaceous Monteith-Gething formations. West of this step, during Laramide compression, a pair of backthrusts nucleated on either side of a minor east-west trending Carboniferous fault and propagated across it in an en échelon pattern. One backthrust ramped laterally across the area and separated the Pink Mountain and Spruce Mountain structures, which both are contained within a 30+ km long pop-up structure above the Besa River Formation detachment.

Glomerspirella fossils confirm the existence of the Upper Jurassic Upper Fernie Formation and Upper Jurassic to Lower Cretaceous Monteith Formation at Pink Mountain.

ACKNOWLEDGEMENTS

I would like to extend my deepest thanks and gratitude to my supervisor Deborah Spratt of the Fold-Fault Research Project (University of Calgary), for financial support and allowing me to steer my own course and set my own goals throughout this thesis project. Debbie was always there for her valuable suggestions and some occasional “nudges” to keep me on track towards the completion of my thesis. I would like to thank Mike Cecile and Larry Lane of the Central Foreland NATMAP Project (Geological Survey of Canada). Mike Cecile gave me the Pink Mountain project, field support, and allowed me to work on my thesis while working at the GSC full time. Larry Lane continued the financial and work-time support despite very difficult GSC financial times.

Sincere thanks go to Vlada Avramovic for his hard work re-processing two seismic sections and to both Kelman Technologies and GX Technology for supporting him. Amit Mehra and Lorne Smith of Petro-Canada, and Cecil Keeping of Sigma Explorations Inc. are thanked for providing the raw seismic data for Vlada to re-process. Conoco Canada Ltd., Shell Canada Ltd., Talisman Energy, and Husky Canada also provided seismic data that greatly aided this research and gave this thesis an extra “punch”! These people deserve a great round of applause and thanks for their valuable assistance: Dave McNeil, Denise Then, Phil Lawrence, Gabriella Carelli, J.P. Zonneveld, Barry Richards, and Mike Staniland.

Finally, I would like to thank you Dad, for providing me with the means to complete my undergrad courses, so I qualified for the Master’s! I present to you this thesis, which symbolizes the return on your investment in my education so long ago!

TABLE OF CONTENTS

Approval page	ii
Abstract	iii
Acknowledgements	iv
Table of Contents	v
List of Figures	viii
 CHAPTER 1. INTRODUCTION	
1.1 Structural and Stratigraphic Problems at Pink Mountain	1
1.2 Location and Access to Pink Mountain	3
1.3 Physiography of the Study Area	3
1.4 Economic Significance of the Pink Mountain Area	5
1.5 Previous Major Geological Studies of the Pink Mountain Area	7
1.6 Methods	12
 CHAPTER 2. STRATIGRAPHY OF THE STUDY AREA	
2.1 Generalized Stratigraphic Summary	15
2.2 Constraints for the Older Geological Units	15
2.3 Proterozoic (Upper Proterozoic)	18
2.4 Cambrian (Lower to Upper Cambrian)	18
2.5 Shelf vs. Platform Stratigraphy of the Lower Ordovician to Lower Devonian	19
2.6 Ordovician	19
2.6.1 Kechika Group (Tremadocian - Arenigian)	19
2.6.2 Skoki Formation (Arenigian - Darrwilian)	21
2.6.3 Beaverfoot Formation (Caradocian - Ashgillian)	21
2.7 Silurian	22
2.7.1 Unnamed Silurian Limestone (Llandoveryian)	22
2.7.2 Nonda Formation (Llandoveryian - Pridolian)	22
2.8 Silurian - Devonian	22
2.8.1 Muncho-McConnell Formation (Pridolian - Lochkovian)	22
2.9 Devonian	23
2.9.1 Lateral Variations of the Lower Devonian	23
2.9.2 Wokkpash Formation (Pragian)	23
2.9.3 Stone Formation (Emsian - Eifelian)	25
2.9.4 Dunedin Formation (Eifelian - Givetian)	25
2.10 Devonian - Carboniferous	27
2.10.1 Besa River Formation	27
2.11 Carboniferous	29
2.11.1 Prophet Formation (Tournasian - Visean)	29
2.11.2 Stoddart Group (Visean - Serpukhovian)	31

2.12 Carboniferous Equivalents to the Prophet Formation	31
2.12.1 Banff Formation (Early - Middle Tournaisian)	32
2.12.2 Pekisko Formation (Middle Tournaisian)	32
2.12.3 Shunda Formation (Late Tournaisian)	32
2.12.4 Debolt Formation (Middle - Late Viséan)	33
2.13 Complications Associated with the Pekisko, Shunda and Debolt Formations	33
2.14 Summary of the Carboniferous and Older Stratigraphy	34
2.15 Permian	35
2.15.1 Belloy or Fantasque Formation (Lower - Upper Permian)	35
2.16 Triassic	35
2.16.1 Montney Formation (Anisian)	35
2.16.2 Doig Formation (Middle Anisian - Ladinian)	35
2.16.3 Halfway Formation (Ladinian)	36
2.16.4 Charlie Lake Formation (Ladinian to Carnian)	36
2.16.5 Baldonnel Formation (Carnian)	36
2.16.6 Pardonet Formation (Norian)	37
2.17 Jurassic	37
2.17.1 Nordegg Member (Pliensbachian - Early Toarcian)	37
2.17.2 Upper Fernie Formation (Kimmeridgian - Lower Volgian)	38
2.18 Jurassic-Lower Cretaceous	38
2.18.1 Monteith Formation (Kimmeridgian - Berriasian)	38
2.19 Cretaceous	45
2.19.1 Cadomin Formation (Valanginian - Hauterivian?)	45
2.19.2 Gething Formation (Hauterivian - Albian)	46
2.19.3 Field Mapping of the Monteith - Gething formations	47
2.19.4 Buckinghorse Formation (Early - Middle Albian)	48
2.19.5 Sikanni Formation (Late Albian - Early Cenomanian)	50
2.19.6 Sully Formation (Late Albian)	50
2.19.7 Dunvegan Formation (Cenomanian)	50
 CHAPTER 3. SUBSURFACE WELL-LOG INTERPRETATION	
3.1 Generalized Drilling Summary in the Trutch-Halfway River Area	51
3.2 Well-Log Interpretation and Methods	51
3.2.1 Data Input	51
3.2.2 Well-Log Interpretation	53
3.3 Previous Well-Log Research in the Halfway River Area	55
 CHAPTER 4. STRUCTURAL GEOLOGY OF THE TRUTCH AND HALFWAY RIVER AREAS	
4.1 Generalized Structural Setting of the Cordilleran Deformation	58
4.2 Formation and Configuration of Thrust Fault Systems	58
4.3 Structural Geology of the Trutch (94G) and Halfway River (94B) Areas	61
4.4 Subsurface Seismic Resolution in the Trutch Area	63

4.5 Surface Structural Geology of the Pink Mountain Anticline	65
4.6 Subsurface Structural Interpretation of Pink Mountain	67
4.7 Surface Identification of the Pink Mountain Backthrust Faults	69
4.8 Stereonet Analysis of Pink and Spruce Mountains	69
4.9 Deep Structural Interpretation of the Pink Mountain Area	71
4.10 Balanced Cross-sections of the Pink and Spruce Mountain Anticlines	72
CHAPTER 5. TECTONIC HISTORY OF THE TRUTCH AND HALFWAY RIVER AREAS	
5.1 Pink Mountain Area in Relation to the Peace River Arch	74
5.2 Precambrian to Cambrian Tectonic History	74
5.3 Ordovician and Silurian Quiescence	77
5.4 Devonian and Carboniferous Tectonism and the Dawson Creek Graben Complex	77
5.4.1 Thickness Variations within the Stoddart Group in the Peace River Area	78
5.4.2 Stoddart Group Depositional Style in the Pink Mountain Area	81
5.4.3 Evidence for Major and Minor Subsurface Normal Faults at Pink Mountain	83
5.4.4 Late Carboniferous to Permian Paleosurface	85
5.5 Uncertain Tectonism during the Permian to Triassic	85
5.6 Jurassic to Tertiary Compressional Tectonics	87
5.6.1 Deposition of the Fernie and Minnes Group and the Columbian Orogeny	87
5.6.2 Laramide Orogeny and Timing of Faults at Pink Mountain	88
5.6.3 Changing Style of the Lily Lake Thrust	89
5.6.4 Relationship between Cretaceous and Carboniferous Faults	89
5.6.5 Origin of the Two Bit Creek and Spruce Mountain en échelon Backthrusts	90
5.6.6 Cenomanian to Present Configuration of Pink Mountain	91
5.7 Generalized Synopsis of the Pink Mountain Tectonic History	91
CHAPTER 6. CONCLUSIONS	95
REFERENCES	98

LIST OF FIGURES

Figure 1-1	Location and surrounding physiography of the Pink Mountain Anticline.	2
Figure 1-2	a) View of Pink Mountain from the Alaska Highway, looking West. b) View of a massive rock avalanche on the northeastern side of Pink Mountain, looking southwest.	4
Figure 1-3	a) Geological map, b) balanced cross-sections, and c) strike sections of the Pink Mountain area.	in pocket
Figure 1-4	Geologic map and structure section of the Pink Mountain Area modified from Hage (1944).	8
Figure 1-5	a) Eastern portion of the Halfway River cross-section (after Thompson 1989). b) Eastern portion of the Cameron River structure cross-section (after Cooper 2000). c) Location map and legend.	10
Figure 1-6	Location map of the digital well and seismic data for the study area.	13
Figure 1-7	Pre-stack depth-migrated seismic sections north and south of Pink Mountain.	in pocket
Figure 1-8	Balanced and restored cross-sections north and south of Pink Mountain.	in pocket
Figure 1-9	Well cross-sections. Each digital section was compiled from Geomatix LogM Model Builder.	in pocket
Figure 2-1	Generalized east-west cross-section showing the distribution of major lithostratigraphic units, and the positions of major facies transitions relative to the Halfway River Map area (modified from Thompson 1989).	16
Figure 2-2	Stratigraphy of the study area (modified from Stockmal 1999, Glass 1997, Pyle and Barnes 2000, and Stott 1991).	17
Figure 2-3	Lateral facies transition between the Ospika Embayment and the MacDonald Platform (modified from Pyle and Barnes 2000).	20
Figure 2-4	Abrupt Silurian-Devonian lithological changes west of the study area (modified from Ross et al. 1993).	24

Figure 2-5	Non-carbonate well log signature below the top of the Dunedin Fm.	26
Figure 2-6	The changing nature of the Devonian-Carboniferous Besa River Formation from southeast to northwest of Pink Mountain (modified from Kidd 1963).	28
Figure 2-7	Three deeper study area wells outlining the older formation nomenclature picks (formation tops from CANSTRAT logs).	30
Figure 2-8	Location and photomicrographs of the Fernie Formation samples (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).	39
Figure 2-9	Location and photomicrographs of the Monteith Formation samples. Representative Late Jurassic to Barremian, benthic, agglutinated foraminifera from sample SHJ-131. Probable age, Oxfordian to Lower Volgian. (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).	41
Figure 2-10	Generalized stratigraphic section of Triassic to Jurassic Strata located on the western flank of the Pink Mountain Anticline (see Figure 2-9 for section location).	42
Figure 2-11	Section 1 outlining the lateral extent of the Monteith Formation, northwest of the Pink Mountain Anticline (see legend on Figure 2-12; modified from Stott, 1998).	43
Figure 2-12	Section 2 outlining the lateral extent of the Monteith Formation, south of the Pink Mountain Anticline (see Figure 2-11 for the section location; modified after Stott, 1998).	44
Figure 2-13	Location and photomicrographs of the three Buckinghorse Formation samples (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).	49
Figure 3-1	Location map for the interpretation wells (green dots).	52
Figure 3-2	The subsurface well-log stratigraphy and synthetic seismogram.	54

Figure 3-3	Carboniferous Dawson Creek Graben Complex and the Stoddart Group isopach map of the Peace River area (modified from Barclay et al. 1990).	56
Figure 4-1	Evolution of a Cordilleran thrust fault system. This diagram is derived from Cordilleran examples assuming plane strain and kink folding (S-fault slip, B - fault angle, modified from Boyer and Elliott 1982).	60
Figure 4-2	View of Pink Mountain looking north. The steeper limb of the Triassic fold is outlined in white (photo by Andrew Miall).	62
Figure 4-3	Location map of the digital well and seismic data for the study area.	64
Figure 4-4	Map of Pink and Spruce Mountains at Halfway River showing stereoplots in the three main fold domains.	66
Figure 4-5	Gamma Ray well-log interpretation of two selected wells on Pink Mountain.	68
Figure 4-6	Composite airphoto mosaic of Pink Mountain from photos taken in 1945. The Two Bit Creek Fault can be seen in the dashed outline (Department of Energy Mines and Resources Canada).	70
Figure 5-1	a) Structure contour map and b) cross-section of the Peace River Arch Precambrian surface (modified from Trotter 1989).	75
Figure 5-2	Basement domains of western Alberta and northeastern British Columbia (modified from Ross et al. 1993).	76
Figure 5-3	Carboniferous-Permian geological and tectonic elements in northeastern British Columbia and Alberta (modified from Barclay et al. 1990).	79
Figure 5-4	Schematic diagram of a) the Carboniferous-Permian Dawson Creek Graben Complex and b) the depositional history of the Stoddart Group (modified from Barclay et al. 1990; see Figure 2-2 for stratigraphy).	80
Figure 5-5	Isopach maps (contours in metres) of the Carboniferous Golata and Kiskatinaw formations in the Trutch and Halfway map areas.	82

Figure 5-6	North-South seismic section showing the Pink Mountain and Elbow Creek Faults (seismic section donated by Petro-Canada).	84
Figure 5-7	Isopach maps (contours in metres) of the Lower Triassic Lower Montney and Upper Montney formations in the Trutch and Halfway map areas.	86
Figure 5-8	Tectonic history of the Pink Mountain area; Carboniferous to Lower Cretaceous (Aptian).	92
Figure 5-9	Tectonic history of the Pink Mountain area; Cretaceous (Albian) to present.	93

1. INTRODUCTION

1.1 Structural and Stratigraphic Problems at Pink Mountain

Figure 1-1 best portrays the immediate question that arises when viewing a topographic map of the Pink Mountain area in northeastern British Columbia: Why is there an isolated high in the Plains east of the Rocky Mountain Foothills?

On earlier (Hage 1944, Taylor 1979) geology maps of Pink Mountain, the rock units are older on the mountain than in the surrounding valleys. There is no surface evidence on or surrounding Pink Mountain of the processes that would locally a) expose older stratigraphy normally found in the Foothills, and b) create the asymmetric profile of the box anticline.

On earlier geology maps, Spruce Mountain (Figure 1-1) was assumed to be the southern portion of the Pink Mountain Anticline (Taylor 1979, Thompson 1989), suggesting an apparent dextral offset along the Halfway River between the major anticlinal axes of Spruce and Pink Mountains. Is the right-stepping fold axis a result of en échelon folding or the action of a minor east-west strike-slip fault similar to that Cecile (1997) proposed for the Marion Lake map area? Are Spruce Mountain and Pink Mountain parts of the same fold? Are the Pink Mountain Anticline and other structures at the eastern edge of the Rocky Mountain Deformation Front a result of thick or thin-skinned structural deformation?

Finally, the precise age and unit nomenclature of the Lower Cretaceous stratigraphy on Pink Mountain and surrounding areas were previously unknown due to the presence of a) an unconformity between the Jurassic and Cretaceous, and b) the lack of fossil evidence in the strata. How old is the sandstone-dominant stratigraphy on Pink Mountain, what is the proper formation name, and how much vertical section does the Jurassic-Cretaceous unconformity remove?

As part of the Geological Survey of Canada's central Foreland NATMAP project, M. Cecile selected the Pink Mountain Anticline to study because of its unique position in front of the foothills and the easy access to the study area.

Figure 1-1: Location and surrounding physiography of the Pink Mountain Anticline. Contour interval is 500 feet, elevation range is 2500 - 6500 feet.

1.2 Location and Access to Pink Mountain

Pink Mountain is located approximately 20 km west of the small town of Pink Mountain at Mile 143 of the Alaska Highway in northeastern British Columbia (approximately 57° N latitude, 123°W longitude). The study area falls within the Trutch (94G) and Halfway River (94B) 1:250,000 map sheets. Motor vehicle access to the northern portion of Pink Mountain is made possible by two gravel roads connected to the Alaska Highway (Figure 1-1). One road leads to a communications tower at the summit of Pink Mountain. The study area is approximately 181 km north of the major town of Fort St. John. Access to the southern portion of the Pink Mountain Anticline on Spruce Mountain, south of the Halfway River, can only be achieved by helicopter and just two limited field traverses were carried out in that area.

1.3 Physiography of the Study Area

Pink Mountain is situated at the western edge of the relatively flat Interior Plains just east of the Rocky Mountain Foothills (Figure 1-1). The elevation of Pink Mountain is 1780 metres at the summit just north of the Halfway River. Pink Mountain appears from the highway as a gently rounded elongate structure that is bisected by two drainages, a branch of Quarter Creek to the north and the Halfway River in the south (Figure 1-1, Figure 1-2a). Local inhabitants of the nearby town of Pink Mountain and industry personnel refer to the three portions as: Little Pink (the northern tip), Pink Mountain, and Spruce Mountain (south of Halfway River).

The name of Pink Mountain is derived from the pink appearance of the exposed bedrock on the mountain during sunset. The normally light brown-grey sandstones appear pink in fading light because of sand grain staining through oxidation of the constituent microscopic iron minerals. In some areas along the west slope, small portions of the sandstones approach a deep red, suggesting a higher concentration of interstitial iron minerals.

Pink Mountain is steeper to the west than the east. The western slope of the mountain approaches 30° and is cut by numerous gullies that are difficult to navigate, but provide the best bedrock exposure of the mountain. The eastern flank of the mountain

Figure 1-2: a) View of Pink Mountain from the Alaska Highway, looking west.

b) View of a massive rock avalanche on the northeastern side of Pink Mountain, looking southwest.

slopes less than 20° and has fewer gullies throughout. The eastern slope has a more fractured, rubbly surface with less “in place” bedrock exposure. Pink Mountain appears symmetric only at the northern tip of Little Pink.

The Interior Plains physiography of the study area is controlled mostly by the bedrock lithology. The more resistant sandstone units form hills or plateaus whereas the more erodable shale units form the sometimes marshy river valleys between Pink Mountain and the low plateaus to the east and the higher elevation foothills to the west (Figure 1-1). In general, deciduous trees and bushes, such as poplars and alder brush grow in the lower elevation areas. Mostly silver birch and aspen trees grow in some of the lower elevation areas with thicker glacial till cover, such as the lowlands near the Blueberry Ranch at the Halfway River (Figure 1-1). Except along steep river cuts, the lower elevation areas are more difficult to traverse and provide minimal bedrock exposure.

Mostly coniferous spruce and evergreen growth cover the higher elevation areas up to the tree line. Above tree line (approximately 1500 metres), discontinuous grasslands with numerous species of small northern flowers characterize the mountain summits along the eastern Foothills. The mostly sandstone rocks of the eastern Foothills and Pink Mountain summits form a bouldery surface from water erosion. These boulder surfaces are difficult to navigate in some places and can complicate seismic exploration (C. Calhoun pers. comm.).

Several rock avalanches occur along the flanks of Pink Mountain. These rockfalls are thought to result from water erosion and freezing of fractures and joints along the steeply dipping slopes of the mountainside. Three large-scale rock avalanches occur along the northeast flank of Pink Mountain, the southeastern end of Pink Mountain at Halfway River, and the central eastern flank of Spruce Mountain (Figure 1-2b).

1.4 Economic Significance of the Pink Mountain Area

Since the building of the Alaska Highway during the early 1940's, various companies have carried out petroleum, coal, and mineral surveys of the study area. To the east of Pink Mountain, several shallow gas fields such as the Julienne Creek and

Julienne Creek North gas fields were successfully drilled and produced gas from Triassic sandstone and carbonate rocks that were gently folded during the formation of the northern Rocky Mountains.

During 1962, the first exploration well, C-090-C/94G2 was drilled near the summit of Pink Mountain (Lepard et al. 1999). This well successfully produced gas, which led to the eventual drilling of fourteen wells on Pink Mountain in the Elbow Creek Gas field (Province of British Columbia Energy, Mines and Petroleum Resources 1979) between 1962 and 1994. This gas field plus the Grassy and Sikanni fields produce gas from deeper Carboniferous rock units. From the late 1960's to the present, numerous oil companies have shot seismic lines along the eastern Trutch and Halfway River map areas to aid in petroleum exploration. All of this activity has improved access to remote areas that would have been otherwise accessible only by helicopter.

Besides natural gas, coal seams have been observed and mapped on the southern slope near the rock slump of Pink Mountain at Halfway River (Figure 1-3a). One study documents the presence of several thin 10 centimetre to 5 metre thick, medium to high-grade coal seams within the sandstone and shale packages (Guardia, 1971). The coal seams only occur along the southern end of Pink Mountain and are not exposed farther north within the steep gullies. These seams are also absent on the northern slope face of Spruce Mountain, which suggests the mostly thin coal seams on Pink Mountain are local occurrences only, and not economic to mine (Guardia, 1971).

Two mineralized gossan zones have been discovered near the base of the sandstones on the southern portion of Pink Mountain. One along the gravel road near the summit and the other within the rock slump near the Halfway River (Figure 1-3a). The gossan zone in the southern rock slump area has the most extensive alteration affecting approximately 20 metres of stratigraphic section. This alteration is most likely hydrothermal because the rock samples collected indicate mineral replacement on a molecular level. One sample, SJH-065, is silica-altered limestone, which has a preserved ammonite fossil only observed in limestone horizons older than the sandstones. Once again, no other gossan zone occurrences have been observed on Pink Mountain, suggesting the limited lateral coverage, and hence, uneconomic value of these deposits.

1.5 Previous Major Geological Studies of the Pink Mountain Area

Shortly after the construction of the Alaska Highway in the early 1940's, C.O. Hage (1944) of the Geological Survey of Canada conducted geological reconnaissance mapping between Fort St. John and Fort Nelson along the Alaska Highway. One of his stops was Pink Mountain and he mapped it as a single continuous anticline from the Halfway to Sikanni Chief Rivers (Figure 1-4). Hage (1944) also documented the presence of minor folds along the eastern limb of the major fold structure he termed the Pink Mountain Anticline (Hage 1944; McLearn and Kindle 1950). An interesting feature on his map is the southern extrapolation of a shallow normal fault along the eastern limb of the Pink Mountain Anticline, which was only observed at the Sikanni Chief River and a small creek two kilometres south of the Sikanni Chief River (Figure 1-4).

Hage (1944) indicates in his cross-section that the Pink Mountain Anticline is bounded by two faults on either side of the fold at the Sikanni Chief River. At Pink Mountain, the anticline is bounded only by an eastern thrust fault (Figure 1-4). Hage (1944) estimates the displacement on the thrust fault to be approximately 350 metres. The most important structural observation on his cross-section is an eastward dipping thrust fault to the east of Pink Mountain Anticline (Figure 1-4). Today, this easternmost fault is observed on several seismic sections that cross the Pink Mountain Anticline.

Using the stratigraphic nomenclature of the 1940's and the Triassic fossils collected, Hage (1944) mapped the rock units of the Pink Mountain area and measured a stratigraphic section along the Sikanni Chief River (Figure 1-4). The oldest Triassic undivided rock units are located in the core and deeper erosional gullies of the Pink Mountain Anticline whereas the outer limbs contain the Lower Cretaceous Bullhead Group sandstones and the Buckinghorse Formation shales. Hage (1944) also recommended that a coal feasibility study be done on Pink Mountain.

D.F. Stott, B.R. Pelletier and D. Gibson, also of the Geological Survey of Canada, carried out the next major mapping and stratigraphic studies in the Trutch area during the early 1960's, which had implications for the Triassic to Cretaceous stratigraphy on Pink Mountain (Pelletier and Stott 1963). Using a more refined stratigraphic column derived from adjacent areas, they concluded that an erosional unconformity removed the upper

Figure 1-4: Geologic map and structure section of the Pink Mountain Area, after Hage (1944).

Jurassic and some of the Lower Cretaceous stratigraphy including the Minnes Group along the eastern Foothills. Therefore the Lower Cretaceous sandstone dominant stratigraphy on Pink Mountain belonged only to the Gething Formation.

R. Thompson (1989) of the Geological Survey of Canada carried out the next major study in the Halfway River (94B) map area in the late 1970's. This work was a structural and stratigraphic refinement of an earlier mapping project conducted by E. Irish (1970) of the Geological Survey of Canada. The main structural conclusions to come out of Thompson's (1989) report are the effect of "blind" subsurface thrusts on the Devonian to Cretaceous stratigraphy and the role of the thick shale sequences of the Besa River Formation as a décollement horizon for the structural deformation (Figure 1-5a).

Within the Trutch and Halfway River map areas, Thompson (1989) proposes that most of the subsurface faulting in the eastern Foothills is thin-skinned and occurs within the Devonian Carboniferous strata above the Besa River Formation. The effects of the blind thrusting are broad elongate box fold structures within the younger strata above the décollement. The cross-sections of Thompson (1989) are among the first published cross-sections to display the box fold structures observed in the field.

The eastern edge of cross-section A-A' in Figure 1-5a is west of Spruce Mountain and displays no structural deformation. However, on the cross-section, the easternmost anticline above the tip of a blind thrust in Devonian-Carboniferous strata is asymmetric with a steeper west limb (Thompson 1989). This structure matches the symmetry of the Pink Mountain Anticline near the edge of the Rocky Mountain deformation front. All of Thompson's (1989) cross-sections are balanced and were constructed without the aid of seismic data.

In his report, Thompson (1989) also analyzes the structural configuration of the Halfway River folds and summarizes the stratigraphy and tectonic evolution of the area. Another major hypothesis is the possible influence on present-day surface structures of pre-existing faults formed during older, separate tectonic events. These deeper faults are the root of the deflected fold axes mapped at the surface within the Halfway River map area and may have deflected the fold axes of younger structures during deformation.

Figure 1-5: a) Eastern portion of the Halfway River cross-section (after Thompson 1989).
 b) Eastern portion of the Cameron River structure cross-section (after Cooper 2000).
 c) Location map and legend.

Unfortunately, Thompson (pers. comm., 2000) could not reinforce his findings with seismic evidence during the time of his study.

A more recent study in the Halfway River area by M. Cooper (2000) of EnCana portrays a different thick-skinned deformation style along the edge of the Rocky Mountain Deformation Front (Figure 1-5b). Using partial seismic coverage along the cross-section as a guide, Cooper (2000) hypothesizes the re-activation of pre-existing deeper faults during the more recent Laramide Orogeny. These faults propagate to the surface and produce broad wavelength, inversion anticlines along the eastern edge of the deformational front.

During the field season of 1996, M. Cecile (1997) of the Geological Survey of Canada conducted a geological reconnaissance of the Sikanni Chief River area in the Marion Lake (94G3) area west of Pink Mountain. There, he found an apparent north-south lateral discontinuity of the strata across the Sikanni Chief River. Cecile (1997) attributed this discontinuity to a deeper re-activated east-west strike slip fault along the Sikanni Chief River and he hypothesized that similar faults in the western Trutch area could explain the deflection and rapid termination of major fold structures. Subsequent field work resulted in the removal of the Sikanni Chief River stratigraphic lateral discontinuity, but the cause of the other fold deflections could still be deeper east-west trending faults (M. Cecile, pers. comm.).

At present, the Geological Survey of Canada is conducting a new multi-disciplinary mapping project termed the Central Foreland NATMAP project (with L. Lane as the chief geologist) in the Trutch and Liard basin areas. In the Trutch area, the main goals of the project are to update contacts drawn from older research, subdivide formations and groups, build cross-sections, and distribute their findings and maps to the public in a digital GIS package.

As part of the Central Foreland NATMAP project, J. Dixon (1999) of the Geological Survey of Canada interpreted the Triassic stratigraphy of over 700 subsurface well logs in the Trutch area. Dixon's (1999) isopach maps indicate dramatic lateral thickness variations within the older Triassic units. Of particular interest are the east-to-west thickness variations of the older Triassic units in the Pink Mountain area.

Also as part of the Central Foreland NATMAP Project, J. White sampled over three hundred locations of the lower Cretaceous for palynological analysis. These samples revealed no preserved pollen due to possible hydrothermal leaching. This leaching could be related to the occurrence of the gossan zones mentioned previously (J. White pers. comm.). Preservation of Jurassic to lower Cretaceous flora and fauna in the Pink Mountain area is poor due to the high thermal maturity (McNeil 2000, J. White pers. comm.). To overcome this problem, Denise Then of GSC Calgary has carefully processed the paleontological samples collected by the author within this interval (Then and Dougherty 1983).

West of the study area, various geologists under M. Cecile and L. Lane have provided more detailed surface geological and structural compilations of the Marion Lake (94/G3), Mount McCusker (94/G4) and Redfern Lake (94/G5) map areas (Cecile 1999, Cecile and Legun 2001, Lane 2001). At present, the printed open file maps are complete and the work on the cross-sections is in progress.

The studies mentioned above provide an excellent background and additional data for the geological and structural evaluation of the Pink Mountain Anticline.

1.6 Methods

In all the published studies researched, there is no conclusive subsurface analysis of the Trutch area that can provide insights into solving the above structural and stratigraphic questions concerning Pink Mountain. Private industry has conducted confidential shallow subsurface surveys on limited areas at or surrounding Pink Mountain, but no lateral relationships between these areas and the overall structural geology of the Rocky Mountain Foothills have been identified. Figure 1-6 outlines the extensive digital seismic and well data acquired from private industry for analysis in this project. This dataset is extensive and some would argue beyond the scope of a Master's thesis. However, analysis of the best quality seismic data provides constraints on the subsurface stratigraphy and structures of Pink Mountain and their relation to the Foothills structural and stratigraphic configuration.

Figure 1-6: Location map of the digital well and seismic data in the study area.

Using the excellent digital dataset, the first step was to map and standardize the subsurface stratigraphy in the digital well logs and then determine the stratigraphic signature in the seismic sections using synthetic seismograms based on sonic logs. To gain better resolution of the deeper seismic units, Vlada Avramovic of the Fold and Fault Research Project has assisted in the geological and numerical re-processing of selected seismic lines north and south of the Pink Mountain Anticline (Figure 1-7). Once the stratigraphy was known and traced on the enhanced seismic sections, the structural geology could be determined, and balanced cross-sections drawn and restored (Figure 1-8). Tectonic controls on sedimentation and erosion patterns were made evident with the construction of well-log cross-sections (Figure 1-9) and isopach maps, and the final results are presented in a three dimensional perspective.

A better understanding of the three-dimensional nature of these structures is extremely important. The recent discovery of natural gas reservoirs within the Pink Mountain Anticline suggests the potential of future discoveries in other adjacent subsurface structures similar in nature to the Pink Mountain Anticline east of the Foothills.

2. STRATIGRAPHY OF THE STUDY AREA

2.1 Generalized Stratigraphic Summary

In his study of the Halfway River (94B) map sheet just south of Pink Mountain, Thompson (1989) summarized the stratigraphy of the area as being deposited on a thinned continental crust of a passive Atlantic-type continental margin. This miogeoclinal continental prism of deposition extended from the Late Proterozoic to the Early Tertiary. In general, Thompson (1989) observes most of the sediments thicken towards the west and the Ordovician to Lower Devonian strata thin dramatically or pinch out at the miogeocline-platform boundary (Figure 2-1). However, the Platform-Shelf boundary varies in lateral position from west to east throughout the Ordovician to Lower Triassic periods (Cecile and Norford 1979). Thompson's (1989) summary of the thinned continental crust profile is postulated and not based on seismic sections.

2.2 Constraints for the Older Geological Units

Figure 2-2 portrays the entire surface and subsurface stratigraphy of the study area. At present, the oldest identifiable unit in the well logs is the Silurian-Devonian Muncho-McConnell Formation. Only 14 wells within the southwestern Trutch area penetrate the middle to early Devonian stratigraphy. The poor seismic resolution and limited well depth constraints prevent positive identification of strata older than the Muncho-McConnell Formation in the cross-sections (Figures 1-3, 1-7, 1-8, and 1-9). Stratigraphy older than the Muncho-McConnell Formation is identified in the study area through extrapolation from measured sections, geology maps, and seismic sections farther west (Cecile and Legun 2001, Taylor 1979, Thompson 1989).

As a result of oil exploration in northeastern British Columbia, industry has concentrated on studies of the Triassic and younger geological units with some analysis of the Carboniferous Debolt Formation (Cecile and Norford 1979, Hunt and Ratcliffe 1959). Hence, most of the drilled wells rarely exceed the depth of the upper Debolt (or Prophet) Formation. The limited interest resulted in good to excellent quality seismic processing for the shallower stratigraphy with increasingly poor resolution at depths below 2.5 seconds two-way travel time. Most structural stack seismic lines have high-

Figure 2-1: Generalized east-west cross-section showing the distribution of major lithostratigraphic units, and the positions of major facies transitions relative to the Halfway River Map area (modified from Thompson 1989).

Figure 2-2: Stratigraphy of the study area (modified from Stockmal 1999, Glass 1997, Pyle and Barnes 2000, and Stott 1991).

amplitude but distorted and discontinuous reflectors at depth. Processing of the time-migrated seismic lines concentrated on the shallower stratigraphy, resulting in the deeper units appearing as multiple horizontal reflectors, which are untrustworthy and cannot be correlated laterally.

To the north of the study area along the border of the Trutch (94G) and the Fort Nelson (94J) map sheets (57° latitude), the older units are imaged more clearly but some of the deeper units do not appear to be laterally continuous. A detailed analysis of these lines is beyond the scope of this thesis. For this study, interpretations were made on two seismic lines within the Trutch (94G) and Halfway River (94B) map sheets that were pre- and post-stack depth migrated by Vlada Avramovic of GX Technology (formerly of Kelman Technologies) to resolve the deep stratigraphic resolution problems (Chapter 4).

2.3 Proterozoic (Upper Proterozoic)

Very little is known about the Proterozoic stratigraphy of the Trutch area. On the Halfway and Ware map sheets, the Proterozoic strata are mostly slate, dolostone, and limestone sequences of approximately Ediacaran age and older (Stott 1991). The Proterozoic strata are mostly unnamed except for the schists and minor limestones of the Misinchika Group observed in the Halfway River (94B) map sheet (Glass 1997, Thompson 1989). The exact nature of the contact between the Cambrian and Proterozoic is unknown throughout the Halfway River (94B) and Ware (94F) map areas (Stott 1991, Pugh 1975). The Proterozoic of the Trutch area appears to be layered (Cook 1993).

2.4 Cambrian (Lower to Upper Cambrian)

The dominant lithologies of the Cambrian stratigraphy are cross-bedded quartzites, sandy dolostones, and siltstones, which have been associated with the Lynx Formation (Pyle and Barnes 2000, Pugh 1975). Periodic phases of uplift and subsidence have caused several unconformities between the various unnamed units (Pugh 1975). Other lithologies northwest of the Ospika River include conglomerate, oolitic limestone, and sandstone. These lithologies could be equivalent to the Gog Group (Pyle and Barnes 2000, Pugh 1975). No information on the Cambrian stratigraphy exists near the Pink

Mountain Anticline; therefore the stratigraphy will be termed Cambrian undivided on the deeper seismic sections. The contact between the Cambrian units and the Ordovician Kechika Group has been described as a regional angular unconformity (Pyle and Barnes 2000, Glass 1997, Cecile and Norford 1979, Pugh 1975).

2.5 Shelf vs. Platform Stratigraphy of the Lower Ordovician to Lower Devonian

Figure 2-3 outlines the complex facies transitions and formation nomenclature changes of the Lower Ordovician to Lower Devonian stratigraphy beneath the Stone Formation. Previous studies in the Ospika River area have identified a major basin to shelf transition zone (Cecile and Norford 1979, Thompson 1989). Within the Ospika Embayment to the west, the Lower Ordovician to Lower Devonian Road River Group consists of the Skoki and Ospika formations plus several unnamed siliclastic and carbonate units (Figure 2-3, Pyle and Barnes 2000, Cecile and Norford 1979). The maximum thickness of the Road River Group approaches 1300 metres near the Ospika River, 100 km southwest of Pink Mountain Anticline (Thompson 1989).

East of the Ospika Embayment and within the MacDonald Platform, the Lower Ordovician to Upper Silurian stratigraphy consists of the Ordovician Skoki and Beaverfoot formations, the unnamed Silurian limestone, and the Silurian Nonda Formation (Figure 2-3). The ages of the boundaries between these units are uncertain (Pyle and Barnes 2000). The cross-sections of this thesis do not extend west into the Ospika Embayment, therefore the stratigraphic nomenclature of the MacDonald Platform will be used to help identify the Lower Ordovician to Lower Devonian stratigraphy on the seismic sections beneath Pink Mountain Anticline.

2.6 Ordovician

2.6.1 Kechika Group (Tremadocian - Arenigian)

The Lower Ordovician Kechika Group consists mostly of shales, limestones and limestone conglomerates. The Kechika Group is continuous throughout the Ospika Embayment and the MacDonald Platform (Pyle and Barnes 2000, Cecile and Norford 1979). The thickness of the Kechika Group ranges from 300 metres in the McDame

A. Eastern edge of Ospioka Embayment, Mid-Ordovician (after Cecile and Norford, 1979)
 B. Eastern edge of Ospioka Embayment, Late Ordovician-Early Silurian (after Thompson, 1989)
 Note: teeth point towards shelf

STRATIGRAPHY

Subsystem	AREA	OSPIKA EMBAYMENT (SHELFBREAK)	MACDONALD PLATFORM (PLATFORM-SHELF)
	SERIES/ STAGE		
Lower Devonian			
Upper Silurian	Pridoli	Brown Siltstone Unit	Muncho-McConnell Formation
	Ludlow	Unnamed Breccia Unit	Nonda Formation
	Wenlock	Unnamed Carbonaceous Limestone Unit	Unnamed Silurian Limestone
Lower Silurian	Llandovery	Unnamed Shale and Quartzite Unit	Beaverfoot Formation
Upper Ordovician	Ashgillian		
	Caradocian		
Middle Ordovician	Darriwilian	Ospika Fm.	Skoki Fm.
	Arenigian		
Lower Ordovician	Tremadocian		
Upper Cambrian	Trempealeauan		

Question marks indicate where biostratigraphic resolution is required.

Figure 2-3: Lateral facies transition between the Ospioka Embayment and the MacDonald Platform (modified from Pyle and Barnes 2000).

(104P) map area (260 km from Pink Mountain) to over 1400 metres in the Ware (94F) map sheet (100 km from Pink Mountain) (Pyle and Barnes 2000, Glass 1997). As with the Skoki Formation, the four members of the Kechika Group will appear on seismic sections and cross-sections as one unit (Pyle and Barnes 2000). In the Mount McCusker map area (94G/4) 61 kilometres to the west, the carbonaceous shales of the Kechika Group have strong cleavage (Cecile and Legun 2001).

2.6.2 Skoki Formation (Arenigian - Darrwilian)

The Skoki Formation consists mostly of dolostone and minor limestones (Pyle and Barnes 2000, Thompson 1989) in which various authors have subdivided these units, but for the purposes of this study, the Skoki Formation will be traced on the seismic sections as one unit (Pyle and Barnes 2000, Cecile and Norford 1979). The Skoki Formation ranges in thickness from 62 metres in the southern Alberta Foothills to 1260 metres in the western half of the Halfway River Map area (Pyle and Barnes 2000, Glass 1997, Thompson 1989). Near the Ospika River (Figure 2-3), the measured thickness of the Skoki Formation is over 600 metres and the contact with the overlying Beaverfoot Formation is an unconformity (Pyle and Barnes 2000).

2.6.3 Beaverfoot Formation (Caradocian - Ashgillian)

In the Mount McCusker map sheet (94G/4) outcrop localities, the dolostones of the Beaverfoot Formation have mudcracks, hematite nodules, and breccias (Cecile and Legun 2001). The thickness of the Beaverfoot Formation dolostones and sandstones ranges from 320 metres near the Ospika River to 500 metres at Pedley Pass 60 kilometres farther east (Pyle and Barnes 2000, Glass 1997). The contact between the Beaverfoot Formation and the above unnamed Silurian Limestone unit is observed as abrupt (Pyle and Barnes 2000).

2.7 Silurian

2.7.1 Unnamed Silurian Limestone (Llandoveryian)

The unnamed Silurian Limestone unit consists of limestones (Cecile and Legun 2001) that are more argillaceous and thin bedded near the base and become more massive upward (Pyle and Barnes 2000). The unnamed Silurian Limestone unit has a total measured thickness of 220 metres near the Ospika River (Pyle and Barnes 2000). The lateral thickness and continuity of this unit is unknown at present and hence, the extent of the unconformable contact with the overlying Nonda Formation is also uncertain (Pyle and Barnes 2000).

2.7.2 Nonda Formation (Llandoveryian - Pridolian)

A characteristic feature of the Silurian Nonda Formation is the presence of chert nodules within the mainly siliceous dolostone beds. The base of the Nonda Formation usually has thin layers of quartzite, quartz sandstone, and limestone (Stott 1991, Glass 1997). At the type section near the Toad River Bridge on the Alaska Highway at mile 318 (210 kilometres northwest of Pink Mountain), the Nonda Formation is 60 metres thick and thickens to 612 metres at Gibault Creek, just west of Fort St. John (Glass 1997). Within the area of the Figure 2-3 MacDonald Platform sections, the mostly massive dolostones of the Nonda Formation were measured at 200 metres thickness (Pyle and Barnes 2000).

2.8 Silurian-Devonian

2.8.1 Muncho-McConnell Formation (Pridolian - Lochkovian)

The lowermost recognizable lithologies on the well logs are the carbonates and minor siliclastics of the Silurian-Devonian Muncho-McConnell Formation (Pyle and Barnes 2000). The generalized description of the Muncho-McConnell Formation lithology is a unit comprising mostly medium to dark grey dolostones with minor siltstone interbeds and shaley horizons throughout (Stott 1991, Glass 1997, Taylor and MacKenzie 1970). The Muncho-McConnell Formation varies in thickness from 60 metres in the foothills to 350 metres in the Sentinel Ranges, 50 kilometres farther

northwest (Glass 1997). On the west half of the Mount McCusker map sheet (94G/4), a middle Silurian quartzite marker unit has been identified (Cecile and Legun 2001). This unit does not exceed 50 metres in thickness and will be included within the Muncho-McConnell Formation.

2.9 Devonian

2.9.1 Lateral Variations of the Lower Devonian

Previous studies by various authors west of the Pink Mountain Anticline document a lateral facies transition from carbonate to siliclastic-minor carbonate sequences within the Lower Devonian units near the Trutch (94G) and Halfway River (94B) map border (Thompson 1989, Ross et al. 1993, Barnes and Pyle 2000). The three Devonian units principally affected are the Muncho-McConnell, Stone and Dunedin formations (Figure 2-4). This siliclastic facies within these three dominant carbonate units is theorized to be caused by tectonism resulting in uplift along the eastern limit of the Ospika Embayment within the Halfway and Trutch map areas during the Lower to Middle Devonian (Ross et al. 1993). On both sections of Figure 2-4, the Slave Point Formation has been removed, and the Besa River Formation rests unconformably on top of the Dunedin or Stone formation (Thompson 1989, Ross et al. 1993). The stratigraphic nomenclature is based on outcrop observations (Thompson 1989, Ross et al. 1993).

2.9.2 Wokkpash Formation (Pragian)

A continuous thin package of sandstone to argillaceous dolostones belonging to the Wokkpash Formation conformably overlies the Muncho-McConnell Formation (Taylor and MacKenzie 1970). The Wokkpash Formation does not exceed a thickness of 50 metres but is traceable on the surface on the 1:50,000 Mount McCusker map sheet (94G/4) (Cecile and Legun 2001). On the well logs, the Wokkpash Formation could be confused for the sandy units at the base of the Stone Formation (Ross et al. 1993). For the purposes of creating large scale cross-sections, the thin Wokkpash Formation is included with the Stone Formation.

Figure 2-4: Abrupt Silurian-Devonian lithological changes west of the study area (modified from Ross et al. 1993).

2.9.3 Stone Formation (Emsian - Eifelian)

Assuming that the carbonates immediately above the Wokkpash or Muncho-McConnell formations belong to the Dunedin Formation, suggests that the carbonates and minor anhydrites that resemble the Muskeg Formation are actually part of the Stone Formation in the study area. However, there is no documentation of anhydrite within the Stone Formation (Glass 1997, Taylor and MacKenzie 1970). The Stone Formation is composed mostly of light grey dolostones and dolostone breccias. The thickness of the Stone Formation increases southwestward from 140 metres at Mount Bertha to 590 metres at the Sentinel Ranges (Taylor and MacKenzie 1970), 260 to 210 kilometres northwest of Pink Mountain respectively.

2.9.4 Dunedin Formation (Eifelian - Givetian)

The Dunedin Formation is composed mostly of dark grey, argillaceous to siliceous limestones with periodic dolomitic horizons. The total thickness of the Dunedin Formation ranges from 240 metres near the town of Pink Mountain to 385 metres at the Sentinel Ranges, 210 kilometres northwest (Taylor and MacKenzie 1970). Well D-87-I / 94G14 has a significant sandstone signature below the base of Dunedin Formation (Figure 2-5). This well is near Section A of Figure 2-4.

In most of the CANSTRAT logs, a thick package of mostly dolostone and minor anhydrite that fits the description of the Muskeg Formation occurs above the Stone Formation (Law 1955, Belyea and Norris 1962). The anhydrite approaches 25% of the total lithology in some wells within this interval and only the Muskeg Formation has significant amounts of anhydrite (Stott 1991, Belyea and Norris 1962). However, in well B-82-F / 94G11, the CANSTRAT log documents dolostone only, but the electric log signature is similar to other wells that have anhydrite present within this interval. This presents a dilemma in the stratigraphic analysis because only seven CANSTRAT well logs intersect the Muskeg Formation interval within the study area. Out of the seven logs, only four document the presence of anhydrite (two have less than 15%) within the carbonates. Without well logs, there is no practical method to determine whether anhydrite is present in the remainder of the study area.

Figure 2-5: Non-carbonate well log signature below the base of the Dunedin Fm.

2.10 Devonian - Carboniferous

2.10.1 Besa River Formation

The dominant lithology of the Besa River Formation is shale, but the composition of the shales varies from silty-cherty at the base to more calcareous with minor carbonate beds toward the top (Thompson 1989). An interesting correlative marker within the Besa River Formation is the Exshaw Shale equivalent layer of silty-cherty shale (Bamber et al. 1968). This horizon is observed near the top of the Besa River Formation on almost every deep well log of the study area dataset. Below the Exshaw equivalent are limestone beds that range in thickness from 30 to 300 metres and these beds are equivalent to the Winterburn, Wabamun and Nisku formations (Stott 1991, Glass 1997, Kidd 1963).

Figure 2-6 outlines the complex relationship between the Besa River shales, the Banff Formation and the Wabamun-Winterburn limestones and siltstones. Within the Trutch (94G) map area, the top of the Banff Formation is indistinguishable within the carbonates and the base is marked by the top of the Wabamun Group (Kidd 1963). For the purposes of this project, the Wabamun Group and Winterburn Formation equivalents will be termed the Carbonate Marker. A problem occurs farther northwest where the Carbonate Marker pinches out and the top of the Banff Formation becomes almost indistinguishable from the carbonates of the overlying Prophet or Pekisko Formation (Figure 2-6). The solution is to include the mostly carbonate lithologies of the Banff Formation within the basal portion of the overlying Prophet Formation and mark the top of the first thick shale (formally at the base of the Banff Formation) as the top of the Besa River Formation (Kidd 1963).

As well, the Carbonate Marker should be identified as a lateral facies transition within the shales of the Besa River Formation (Figure 2-6). The importance of this Carbonate Marker within the Besa River Formation will become apparent when analyzing the structural deformation of the Besa River Formation on a large scale. The more resistant limestone and siltstone “wedge” within the Besa River Formation shales to the east of the study area could control the position and initiation of the décollement zone

Figure 2-6: The changing nature of the Devonian-Carboniferous Besa River Formation from southeast to northwest of Pink Mountain (modified from Kidd 1963).

within the Besa River Formation farther west. As well, the more siliceous sequences within the Carbonate Marker could be structural hydrocarbon traps, provided the temperatures are not outside the oil or gas-window stage of hydrocarbon development.

2.11 Carboniferous

2.11.1 Prophet Formation (Tournasian - Viséan)

Within northeastern British Columbia, the subsurface Devonian to Carboniferous stratigraphy is complicated by incorrect nomenclature assignments due to lateral facies changes and unconformities. The thesis study area has a combination of northern and southern formation names that range from the district of Mackenzie to the southern Alberta Foothills. Most often, the formation names assigned to the CANSTRAT well logs are loosely based on stratigraphic type sections found hundreds of kilometres south of the study area (Bamber and Mamet 1978, Belyea and Norris 1962). In numerous wells, the formation tops for the Debolt (and its subdivisions), Shunda, Pekisko, and Banff formations usually fall on gamma and sonic log intervals that are not consistent and cannot be reliably traced laterally (Figure 2-7).

The type section of the Prophet Formation is along Bat Creek (80 km northwest of Pink Mountain) within the Trutch map area (Bamber and Mamet, 1978). The Prophet Formation was mapped at the surface along the western halves of the 1:250,000 scale Trutch and Halfway River map sheets (Irish 1970, Taylor et al. 1999, Taylor 1979). In the subsurface, the eastern halves of these map sheets retain the older unit names based mostly on southern Alberta stratigraphy (Irish 1963). The Prophet Formation is composed mostly of limestone, dolostone and chert, and the formation can be subdivided into three members, from oldest to youngest: A, B, and C.

Member A of the Prophet Formation contains a great variety of limestone classifications and compositions such as radiolarites and argillaceous wackestones. The thicker shales at the base of Member A are less calcareous and form a gradational contact with the underlying Besa River Formation. In some studies, the base of the Prophet Formation extends down close to the base of the Pekisko Formation (Bamber and Mamet 1978, Bamber et al. 1968).

Figure 2-7: Three deeper study area wells outlining the older formation nomenclature picks (formation tops from CANSTRAT logs).

Member B is composed mostly of siliceous limestones and dolostones with interbedded chert. The chert horizons distinguish Member B from the other members. The limestones and wackestones of Member C occur at the top of the Prophet Formation as a thin horizon that is difficult to identify on the well logs and is included in Member B in this study (Bamber and Mamet 1978, Irish 1963).

2.11.2 Stoddart Group (Visean - Serpukhovian)

Above the Prophet Formation, the Stoddart Group consists mostly of sandstone and shale with rare carbonate beds. The type section is in a subsurface well drilled near Fort St. John (Bamber and Mamet 1978). The Stoddart Group can be subdivided into, from oldest to youngest, the Golata, Kiskatinaw and Taylor Flats formations (Bamber and Mamet 1978, Irish 1963). The thickness of the Stoddart Group varies from zero metres to over 400 metres within the study area (Barclay et al. 1990).

The Golata Formation consists mostly of shales and interbedded argillaceous limestones and sandstones (Bamber and Mamet 1978). The contact between the Prophet and Golata formations form a gamma ray and sonic log peak that is distinctive and can be traced laterally (Figure 2-7). When present in the study area, the top of the quartz sandstones and dark grey shales of the Kiskatinaw Formation is marked by an unconformity (Barclay et al. 1990, Bamber and Mamet 1978). The predominantly limestone and dolostone beds of the Taylor Flats Formation are not observed in the subsurface CANSTRAT logs of the study area. This could be the result of a lateral facies transition to the northwest or the sub-Triassic unconformity (Bamber and Mamet 1978).

2.12 Carboniferous Equivalents to the Prophet Formation

Within the Trutch map area, the eastern subsurface well log interpretations mainly use the Banff, Pekisko, Shunda, and Debolt formations as their major formation picks for the equivalents to the Prophet Formation (Irish 1963). All of these formations have their respective type sections located south of the Peace River area in Alberta. For example, the Banff Formation type section is located in southern Alberta at Mt. Rundle (Bamber

and Mamet, 1978). As documented below, the lithologies of the 14 deep wells in the study area are much different than those of the type sections in Alberta.

2.12.1 Banff Formation (Early - Middle Tournaisian)

Above the Carbonate Marker of the Besa River Formation are the Banff Formation shales and minor carbonates. In the Trutch (94G) and Halfway River areas (94B), the thickness of the Banff Formation ranges from 244 metres and thickens westward to 381 metres (Bamber and Mamet 1978). Some of the CANSTRAT formation picks (Well C-037-L / 94G/07) for the Banff Formation include over 50 percent carbonate lithologies within the shale packages. This is a substantial difference from the Banff Formation type section (Bamber and Mamet 1978). However, recent studies segregate the Banff Formation into six members from A to F. Members C and D are present in northeastern British Columbia and they consist mostly of carbonates (Glass 1997; B. Richards, pers comm.).

2.12.2 Pekisko Formation (Middle Tournaisian)

The overlying Pekisko Formation is composed mostly of skeletal limestone and wackestone with minor shale and argillaceous limestone (Bamber and Mamet, 1978). A distinctive feature of this formation is the presence of crinoids within this interval as logged by CANSTRAT services. The thickness of the Pekisko Formation is usually less than 72 metres. Some intervals mapped as the Pekisko Formation (Well C-037-L / 94G/07 - above the Banff Formation) are composed mostly of dolostone, a lithology not associated with the type section of this formation (Bamber and Mamet, 1978).

2.12.3 Shunda Formation (Late Tournaisian)

In most studies, the Shunda Formation limestones, and minor calcareous shales and dolostones occur directly above the Pekisko Formation. Throughout the study area, the thickness of the Shunda Formation varies from 72 to 183 metres (Law 1981, Bamber and Mamet, 1978). Various private industry personnel describe the sonic and gamma ray

log signatures of the Pekisko and Shunda formations as very difficult to determine in the best of conditions (Figure 2-7; H. Plint pers. comm.).

2.12.4 Debolt Formation (Middle - Late Viséan)

This package of rocks is known as the Debolt Formation in wells drilled into the platform in the eastern part of the study area, but as the Prophet Formation in outcrops west of Pink Mountain, so it is labeled Debolt in well data and Prophet in maps and cross-sections. The lower Debolt consists mostly of thick packages of micritic to skeletal limestone with thinner calcareous shale and argillaceous limestone interbeds. The proportions of chert and dolostone increase in the lower Debolt westward towards the Foothills (Bamber and Mamet 1978).

The Upper Debolt Formation consists of limestone, chert, and minor shale packages and in most areas, the contact between the Debolt and overlying Golata Formation is an unconformity, which sometimes removes the upper portion of the Debolt Formation (Law, 1981). The entire Debolt Formation reaches a maximum thickness of 366 metres but thins toward the northwest (Law 1981).

2.13 Complications Associated with the Pekisko, Shunda and Debolt Formations

The Debolt Formation can be subdivided into five members based on the argillaceous and carbonate content of the lithologies: the lower carbonate, the lower argillaceous, the middle carbonate, the upper argillaceous, and the upper carbonate units. (Law, 1981). However, these subdivisions are difficult to place accurately on a well log and are impossible to trace across a large-scale seismic section (Figure 2-7). As seen on the well log of Figure 2-7, there is little contrast in the sonic and gamma ray logs from the top of the Besa River Formation to the upper Debolt Formation. Thus, the lithologies of the Debolt subdivisions must be determined through detailed chip sample analysis of each well, which is beyond the scope of this thesis.

Another complication is the lateral equivalency of the Shunda and Pekisko formations in the Fort St. John area and the lateral equivalency of the Shunda and Besa River formations northwest of the Trutch map sheet (Bamber and Mamet 1978, Bamber

et al. 1968). The obvious question is what lithology percent dictates the nomenclature, and are these transitions visible on larger scale seismic sections? As documented above, it is not feasible to continue with the mostly southern Alberta formation nomenclature in this study area because of: a) the variability of the lithologies between the old formation picks, and b) the inability to distinguish the older formation tops in electric logs and seismic sections (Figure 2-7).

To solve the problems mentioned above, the interval between the Besa River and Golata formations will be designated as part of the Prophet Formation until seismic evidence indicates the lateral facies transition of the Prophet Formation carbonate stratigraphy into the Besa River Formation shale sequences (Bamber et al. 1968). Where possible, the Prophet Formation will be subdivided into its constituent members of A and B-C. For the 14 deep well logs of the study area, the CANSTRAT lithology analysis of the Carboniferous interval fits the criteria for members of the Prophet Formation.

2.14 Summary of the Carboniferous and Older Stratigraphy

A detailed study of the sample chips and core intervals from the deepest wells in the study area would be necessary in order to standardize the stratigraphic units and nomenclature of the subsurface. A subsurface lithological and paleontological study should be combined with a similar analysis of the Cambrian to Devonian stratigraphy that crops out west of the study area. Unfortunately, such a detailed comparison is beyond the scope of this thesis and would constitute a project in itself. Once more information is known, nomenclature of the deeper stratigraphic units in this thesis may change, but their geometries should not.

The configuration of the Ordovician to middle Devonian stratigraphy as seen in seismic data in the region (Figures 1-5b and 1-7) suggests that a) tectonic activity occurred syndepositionally over at least part of this time interval, and b) different structural styles in the Besa River Formation and younger units may be related to changes in geometry and lithology of the underlying rocks. These topics are covered more fully in chapters 4 and 5.

2.15 Permian

2.15.1 Belloy or Fantasque Formation (Lower - Upper Permian)

In some areas within the Trutch map sheet, a discontinuous thin 25 to 75 metre package of cherts, shales, and calcareous sandstones of the Permian Belloy or Fantasque Formation occurs beneath the Montney Formation (Bloy and Scott 1993, Bamber et al. 1968). At the scale of this cross-section, the Belloy Formation is too small to trace on the seismic data and where present, it is included within the underlying formations. In some localities, a glauconitic shale unit near the base can be recognized in the well logs and characterizes the Belloy Formation (Leggett et al. 1993).

2.16 Triassic

2.16.1 Montney Formation (Anisian)

The base of the Montney Formation is an extensive sub-Triassic unconformity and the formations below it vary from Permian to Mississippian in age (Gibson 1971, Pelletier 1964). The unconformity between the Montney Formation and older strata is a marker horizon in well logs (Figure 2-7). The Montney Formation (Grayling Formation equivalent at surface) consists mostly of a lower package of interbedded dark grey siltstones and shales whereas the upper portion is composed of interbedded light brown siltstones and sandstones. The upper and lower portions of the Montney Formation may be discernable on well logs, but this division is not supported by core analysis. The depositional environment of the Montney Formation varied from deep to shallow water during the Early Triassic (Davies 1997, Pelletier, 1964). The thickness of the Montney Formation within the Trutch area is highly variable from 200 to 500 metres.

2.16.2 Doig Formation (Middle Anisian - Ladinian)

The Doig Formation (Toad Formation equivalent at surface) was deposited in a more varied shallow to deeper water environment during the Anisian period of the middle Triassic. The lithologies include shale, siltstone, minor sandstone, and a phosphatic shale layer at the base that is a marker horizon in well logs (Evoy 1997, Davies et al. 1997).

The thickness of the Doig Formation is variable from 365 metres in the Peace River area (Pelletier 1964) to 100-300 metres in the Trutch area (Gibson 1971).

2.16.3 Halfway Formation (Ladinian)

Calcareous sandstones and minor limestones of the Halfway Formation (Liard Formation equivalent at surface) were deposited in a mostly shallow water environment during the Middle Triassic. The maximum thickness of the Halfway Formation is 430 metres near Peace River (Pelletier 1964) and decreases northward to 50-70 metres within the Trutch area (Gibson 1971).

2.16.4 Charlie Lake Formation (Ladinian to Carnian)

The contact between the Charlie Lake Formation and the mostly sandstone lithology of the Halfway Formation is difficult to identify in the field. In outcrops in the Marion Lake map sheet (94G/3), Dr. Mike Cecile (pers. comm.) identified sandstones similar to the Halfway Formation within the base of the Charlie Lake Formation. The lithologies of the Charlie Lake Formation range from massive and reef limestones to sandstones that are interrupted by numerous unconformities. For this study, the Coplin unconformity was identified on the well logs by J.P. Zonneveld of the Geological Survey of Canada, and this is used as the base of the Charlie Lake Formation in the cross-sections. The age of the Charlie Lake is thought to be within the middle to late Triassic, and the thickness ranges between 100 and 150 metres in the Trutch area (Davies et al. 1997, J. Zonneveld pers. comm.). The base of the Charlie Lake Formation is not exposed at Pink Mountain.

2.16.5 Baldonnel Formation (Carnian)

The contact between the Charlie Lake and Baldonnel formations is a subject of debate (M. Orchard and J. Zonneveld pers. comm.). Resistant massive grey limestones that usually form ridges or cliffs characterize the Baldonnel Formation (Gibson 1971). These limestones reach a maximum thickness of 200 metres near the study area. On Pink Mountain, thin 50 centimetre chert horizons have been observed throughout the

Baldonnell Formation. The Baldonnell thins to the north and east of the Trutch area (Davies et al. 1997).

2.16.6 Pardonet Formation (Norian)

The contact between the Pardonet and underlying Baldonnell Formation appears to be gradational at Pink Mountain but is mapped as an unconformity in other surface and subsurface localities within the Trutch area (Johns et al. 2000, Davies et al. 1997, Pelletier 1964). The Pardonet Formation was deposited as a result of a marine transgression (M. Orchard and M. Jones pers. comm.) and a distinguishing feature of the limestones is a brown-weathering platy appearance. The thickness of the Pardonet Formation ranges from a maximum of 610 metres at Peace River (Pelletier 1964) to 150-200 metres at Halfway River (Thompson, 1989). Both micro- and macro-paleontological analysis constrain the age of the Pardonet Formation to be Norian, within the late Triassic (Davies et al. 1997; Pelletier 1964; M. Orchard and M. Johns, pers. comm.). Approximately 20 km north of the Prophet River (65 kilometres northwest of Pink Mountain), the Pardonet Formation is absent (Pelletier 1964).

The Pardonet Formation has abundant thalattosaur fossil “bone beds” in many areas along the western limb of the Pink Mountain Anticline (Nicholls et al. 1998, Storrs 1991). On the western flank of Pink Mountain, both the Baldonnell and Pardonet formations are highly fractured, exhibit some bitumen staining and have a mild to strong petrolific odor.

2.17 Jurassic

2.17.1 Nordegg Member (Pliensbachian - Early Toarcian)

An unconformity separates the Pardonet Formation from the 20 to 60 metre thick package of Jurassic shales belonging to the Fernie Formation (Stott 1998, Gibson 1992). Of the several Fernie Group subdivisions made farther south (Stott 1998), only two exist within the Trutch area, the calcareous marine shales that resemble the Nordegg Member and the marine to near shore facies of the upper Fernie.

Dr. Fabrice Cordey of the University of Lyons, France identified radiolarians of Pliensbachian to early Toarcian (lower Jurassic) age in a sample (SJH-123a) of calcareous shale from the lower Nordegg Member (M. Orchard, pers. comm.). Flattened ammonites from the species *Dumortieria* were found within the same calcareous shales of the Nordegg Member. These ammonites were identified by R. Hall of the University of Calgary and the Pink Mountain ammonites differ from the *Acanthopleuroceras* species of ammonites found by Stott (1998) within the lower Fernie Formation in the Trutch and Halfway River map areas (R. Hall pers. comm.). The nomenclature for this lower calcareous member of the Fernie Formation is a matter of debate (Poulton et al. 1990) but many private industry personnel and scientists refer to this interval as the Nordegg Member (G. Carelli pers. comm.). Until new information becomes available, the lower calcareous member of the Fernie Formation will be referred to as the Nordegg Member.

2.17.2 Upper Fernie Formation (Kimmeridgian - Lower Volgian)

An unconformity exists between the lower calcareous member and non-calcareous shales of the Fernie Formation and this could be one of the reasons for the decrease in thickness of the Fernie Formation north of the Peace River area (Stott 1998).

As for the upper member of the Fernie Formation, Dr. D. McNeil (2000) of the Geological Survey of Canada has identified *Reophax* sp. *Labrospira goodenoughensis* (Chamney 1978) and *Ammobaculites alaskensis* (Tappan 1955) of upper Jurassic age in a marine shale of sample SJH 131. This shale is unconformably overlain by a thick sequence of conglomeratic, coarse-grained sandstones (Figure 2-8).

2.18 Jurassic-Lower Cretaceous

2.18.1 Monteith Formation (Kimmeridgian - Berriasian)

At Pink Mountain, the Monteith Formation consists of mostly fine- to coarse-grained sandstones with thin (less than a metre) shale beds and thick (between 1 and 30 metres) conglomerate packages. Thin coal horizons are sometimes found within the Monteith Formation. The shale horizons have a near-shore to fluvial-deltaic depositional environment (Stott 1998). The conglomerates are mostly fluvial in origin

LOCATION MAP

Saturnella brookeae
(Hedinger)

Labrospira freboldi
(Hedinger)

Labrospira goodenoughensis
(Chamney)

Ammobaculites
(sp. 6 of Wall, 1983)

Recurvoides canningensis
(Tappan)

Ammobaculites alaskensis
(Tappan)

Representative Late Jurassic to Berriasian (probably Oxfordian to Lower Volgian), benthic, agglutinated foraminifera from sample SJH-123. *Ammobaculites alaskensis* and *Ammobaculites* sp. 6 also occur in sample SJH-131.

NOTE: All photomicrograph distances are in microns.

Figure 2-8: Location and photomicrographs of the Fernie Formation samples (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).

(R. MacNaughton, pers. comm.) and the pebbles are mostly light grey chert with minor quartzite. The conglomerate pebbles are well rounded and range in size from 5 millimetres to 3 centimetres. The thickness of the Monteith Formation ranges from 700 metres in the western mountains to 300 metres in the Foothills and finally to zero in the subsurface of the Plains (Stott 1998).

In 1999, the existence of the Monteith Formation at Pink Mountain was confirmed as a result of the discovery of *Glomospirella*, *Saturnella brookeae*, and *Labrespir goodenoughensis* microfossils. These microfossils have an Oxfordian - Kimmeridgian (and possibly Berriasian) age range in the upper Jurassic (Figure 2-9). All specimens were discovered in thin shales in between the thicker sandstone-conglomerate packages of the lower Monteith Formation (Figure 2-10).

The stratigraphic section of Figure 2-10 was compiled from a continuous section of the Pardonet Formation to the Monteith Formation. The most significant observation obtained from the section is the gradational contact between the Fernie and Monteith formations. L. Lane and M. Cecile of the Geological Survey of Canada also observe a gradational contact between the Fernie and Monteith formations just west of the Pink Mountain Anticline (L. Lane and M. Cecile, pers. comm.). Their observations are in agreement with the section of Figure 2-10 on the Pink Mountain Anticline.

On the western flank of the Pink Mountain Anticline, the Monteith Formation exhibits some bitumen staining but at one site the conglomeratic portion of the Monteith (?) Formation has a major concentration of bitumen and coal fragments that appear to have a high thermal maturity (see Figure 2-9 for location). This suggests the possibility of the generation, migration and accumulation of hydrocarbons during some stage of the development of the Pink Mountain Anticline.

In a continuing study initiated in the early sixties, Stott (1998, 1982, and Stott et al. 1963) have mapped the Fernie to Gething formations within the Trutch and Halfway River map sheets. In Stott's (1998) cross-sections north and south of the Pink Mountain Anticline, the Monteith Formation pinches out on section 1 (Figure 2-11), to the northwest of the study area, but is still present in the southern section (Figure 2-12). If the zero edge of the Monteith Formation were drawn between the sections on the Figure 2-11

LOCATION MAP

Labrospira goodenoughensis
(Chamney)

Ammobaculites alaskensis
(Tappan)

NOTE: All photomicrograph distances are in microns.

Figure 2-9: Location and photomicrographs of the Monteith Formation samples. Representative Late Jurassic to Barremian, benthic, agglutinated foraminifera from sample SHJ-129. Probable age, Oxfordian to Lower Volgian (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).

Figure 2-10: Generalized stratigraphic section of Triassic to Jurassic strata located on the western flank of the Pink Mountain Anticline (see Figure 2-9 for section location).

Figure 2-11: Section 1 outlining the lateral extent of the Monteith Formation, northwest of the Pink Mountain Anticline (see legend on Figure 2-12; modified from Stott, 1998).

Figure 2-12: Section 2 outlining the lateral extent of the Monteith Formation, south of the Pink Mountain Anticline (see Figure 2-11 for section location; modified from Stott, 1998).

index map, it would occur just east of, or cross obliquely through the Pink Mountain Anticline.

As a result, the Monteith Formation occurs on the Pink Mountain Anticline as a discontinuous thin unit interrupted by an erosional unconformity overlain by either an intermittent Cadomin Formation or the Gething Formation. The Monteith Formation most probably pinches out toward the northern end of the Pink Mountain Anticline. This would explain why other scientists have concluded that the Monteith Formation is absent within the study area (Gibson 1992, Taylor 1979).

2.19 Lower Cretaceous

2.19.1 Cadomin Formation (Valanginian - Hauterivian?)

Previous studies have not yielded conclusive evidence for the presence of the near-shore to fluvial-deltaic sequences of the Lower Cretaceous Cadomin and the upper Jurassic to Lower Cretaceous Monteith formations in the eastern Trutch (94G) map area of Northeastern BC (Stott 1998, Gibson 1992). Within the southeastern corner of the Marion Lake map sheet, Cecile and Legun (2001) observes a discontinuous 50 metre thick sequence of conglomerates that they call the Cadomin Formation. The well-rounded clasts of the conglomerates are mostly light to medium-grey chert pebbles with minor (less than 10 percent) white to grey quartzite pebbles. These clasts are in a medium to coarse-grained sandstone matrix

This hypothesized Marion Lake Cadomin Formation occurs between the sandstone, shale and conglomerate sequences of what Dr. Cecile (pers. comm.) theorizes as being the Monteith and Gething formations. On the western limb of the Pink Mountain Anticline, several packages of similar chert and quartzite-pebble conglomerates are observed but cannot be correlated laterally. Some of these conglomerates, such as those observed at Halfway River approach 30 metres in thickness and lie directly on top of the upper Fernie Formation.

At the Halfway River locality, geological assistant Scott Carter found shell fragments near the base of the conglomerates, which were analyzed by Terry Poulton and colleagues of the Geological Survey of Canada. Unfortunately, the fragments could only

be identified as possibly four different marine (?) bivalve species, but additional traverses should be done in the Halfway River area to determine if more bivalve fossils can be found in the conglomerate (T. Poulton pers. comm.).

In summary, in the Trutch area the Cadomin Formation may only have been preserved in intermittent channels, that cut down into and may have removed the underlying Monteith Formation and some of the Fernie Formation in places (M. Cecile and C. Schroder-Adams, pers. comm.; Stott 1998). Another problem is that the bivalve fossils indicate a marine environment for the Cadomin Formation. The combined thickness of the Monteith and Gething formations in the study area ranges from 150 to 300 metres. Further south, in the Peace River region, the combined thickness of the two formations and the Cadomin approaches two kilometres (Stott 1998, Gibson 1992).

2.19.2 Gething Formation (Hauterivian - Albian)

The depositional environment of the Gething Formation can be described as near-shore to fluvial-deltaic (Gibson 1992, Stott, 1998). Throughout the Pink Mountain Anticline, the Gething Formation sandstones exhibit prominent channel cross-bedding, fluid escape structures, coaly plant fragment layers and well-developed burrowing horizons. At Pink Mountain, the basal contact of the Gething Formation is an unconformity on top of either the Cadomin or Monteith Formations (Stott 1998, Gibson 1992). On the northern tip of "Little Pink", the uppermost chert and quartzite-pebble conglomerates and finer-grained sandstones of the Gething Formation are overlain in sharp contact with the shales of the Buckinghorse Formation. This non-gradational contact is observed in other areas surrounding Pink Mountain and within Dr. Claudia Schroder-Adams's study area along the Buckinghorse River, 25 kilometres farther north (C. Schroder-Adams, pers. comm.). However, this hiatus is not major, based on the early Albian paleontologic age determined from Sample HQB-39, obtained in this area.

With the exception of unidentifiable plant fragments, preserved fossil evidence in the Gething Formation of the Trutch and Halfway River map areas is scarce (Gibson 1992, Thompson 1989). However, a Haydrasaurus dinosaur footprint found near the summit of Pink Mountain, resembles footprints found in other Gething Formation

localities in the Trutch map area (B. Nicholls pers. comm.). The plant fragments preserved in the sandstone sequences are all coaly, which indicates the Gething Formation may have a high degree of thermal maturity. The high temperature could destroy fossil evidence, especially pollen spores (J. White pers. comm.).

Evidence for thermal alteration of the Pink Mountain strata is found within two gossan zones located along the eastern flank of the anticline at Halfway River and near the summit of Pink Mountain. Here, thermal alteration by migrating hot fluids has affected the Triassic Pardonet to Jurassic-Cretaceous Monteith formations. Movement of hot fluids along faults could have heated the Cretaceous strata enough to destroy the fossils and pollens. Dr. James White of the Geological Survey of Canada was unable to detect pollen within the 300+ samples he collected within the Triassic to Lower Cretaceous shale sequences in the Trutch area (J. White pers. comm.).

The Gething Formation reaches a maximum thickness of 1100 metres just south of Williston Lake (Gibson 1992). The overall age of the Gething Formation is estimated to range from the Aptian to early Albian (Gibson, 1992, Thompson 1989). From field observations along the Pink Mountain Anticline, the Gething Formation thickness is less than 300 metres. At Pink Mountain, previous paleontological results on a sample just below the contact with the Buckinghorse Formation reveal a Cretaceous age with an indeterminate stage (Gibson 1992).

2.19.3 Field Mapping of the Monteith - Gething formations

Field identification of the Monteith Formation in the Trutch map area is complicated by the similar fluvial-deltaic depositional environments of both the Monteith and Gething formations (Stott 1998, Gibson 1992). In outcrop sections just above the Fernie Formation, channel cross-bedding and burrowing horizons similar to those found in the Gething Formation have been observed in the stratigraphic position that should be occupied by the Monteith Formation. After numerous traverses on Pink Mountain, the only weak qualifier to distinguish between the Monteith and Gething formations was the color.

The fresh surfaces of the sandstones, conglomerates and shales appear to change color from a dark grey to dark brown toward the top of the formation. Dr. Claudia Schroder-Adams (pers. comm.) also documented this color difference on her subsequent traverses of the Monteith and Gething Formations along the Sikanni Chief River. There also appear to be more prominent channels and more abundant clean sandstones in the Gething Formation. However, brown and grey units have been identified in both the Monteith and Gething formations (Stott 1998, Gibson 1992). Taylor (1979) and Thompson (1989) proposed that a period of erosion after deposition of the Cadomin Formation removed the Cadomin, Monteith, and some of the Fernie Formation along the eastern half of the Trutch map area. This has resulted in the Gething Formation being deposited on top of the mid- to lower Fernie Formation.

2.19.4 Buckinghorse Formation (Early - Middle Albian)

The Buckinghorse Formation consists of shales and minor iron-rich siltstones that unconformably overlie the Gething Formation. Storm deposits of coarser sediments periodically interrupted the deposition of the Buckinghorse Formation shales in an offshore shelf environment (Schroder-Adams 1999). In general, the Buckinghorse Formation has several subdivisions defined by various authors (Stott 1982), which will not be evaluated in this study. However, a previously undocumented layered unit of coarse-grained sediments has been discovered by the author near the base of the Buckinghorse Formation. This anomalous sequence is a dolostone identified by Dr. Claudia Schroder-Adams (pers. comm.) of Carleton University.

The total thickness of the Buckinghorse Formation is 1128 metres in the C-54-H/094/G-2 well (528971m E, 6328629m N; Stott 1982). In the study area, the total subsurface well thickness measured approaches 975 metres. From previous micropaleontological studies, the Buckinghorse Formation has a late early Albian to middle Albian age within the lower Cretaceous (Schroder-Adams 1999, Stott 1982). The oldest age determined for the Buckinghorse Formation in the study area is early to middle Albian (Figure 2-13), which agrees with previous studies. The micropaleontologic ages

Figure 2-13: Location and photomicrographs of the three Buckinghorse Formation samples (species identifications and age interpretations by D.H. McNeil, Geological Survey of Canada, Calgary).

(McNeil 2000) were obtained from three Buckinghorse Formation samples flanking the Pink Mountain Anticline (Figure 2-13).

2.19.5 Sikanni Formation (Late Albian - Early Cenomanian)

The youngest geological unit exposed at the surface within the Pink Mountain Anticline area is the lower Cretaceous Sikanni Formation. The type section for the Sikanni Formation (at the intersection of the Alaska Highway and the Sikanni Chief River) has four sequences of lower to middle shore-face sandstone separated by silty mudstones with a thickness range of 70 to 115 metres (Stott 1982, Schroder-Adams 1999). The Sikanni Formation is only observed at two locations along the Pink Mountain access road and the major features of this formation are the prominent channels and small bivalve horizons. The basal contact is gradational with the Buckinghorse Formation. Previous paleontologic studies constrain the Sikanni Formation to a late Albian age (Stott 1982).

2.19.6 Sully Formation (Early Cenomanian)

The Sully Formation mostly consists of recessive shales and siltstones that are thought to be conformable with the underlying Sikanni Formation sandstones. The Sully Formation and younger strata do not crop out in the western map sheet (94/G2W) of Pink Mountain but are included in the seismic sections that extend farther east. The maximum thickness of the Sully Formation in the study area is 207 metres (Stott 1982).

2.19.7 Dunvegan Formation (Cenomanian)

The sandstones and conglomerates of the Dunvegan Formation are thought to be conformable with the Sully Formation. The base of the Dunvegan Formation is not well exposed in the southern Trutch area. The thickness of the Dunvegan Formation ranges from 125 to 180 metres throughout the study area (Stott 1982).

3. SUBSURFACE WELL-LOG INTERPRETATION

3.1 Generalized Drilling Summary in the Trutch-Halfway River Area

In the early 1970's to 1980's, industry conducted extensive initial petroleum exploration in the Trutch – Halfway River area (Figure 3-1). This past and on-going exploration has yielded numerous gas reservoir discoveries at various stratigraphic levels throughout the region. The two main gas pay zones occur within the fractured Triassic Charlie Lake to Pardonet formations and the top of the undivided Carboniferous Prophet Formation (Debolt Formation) (Lepard et al. 1999). As a result of the deeper reservoir potential of this area, most of the wells surrounding Pink Mountain have been drilled to the Carboniferous Prophet Formation level, between 800 to 2500 metres depth. Some eastern exploration wells, such as A-50-D-94G/1 and C-04-G-94G/7 have penetrated to the Devonian Stone Formation (Figure 1-9).

3.2 Well-Log Interpretation and Methods

3.2.1 Well-Log Data Input

For higher interpretation accuracy, digital well-logs were obtained for all of the 150 wells that penetrate below the Carboniferous Stoddart Group in the Trutch and Halfway River map areas (Figure 3-1). The LAS format ascii files were obtained from the Geological Survey of Canada (through Datashare Inc.) and from the University of Calgary research agreement with QC Data Inc. Calgary. Ron Stefik of the British Columbia Oil and Gas Commission provided deviation information for selected wells in structural cross-sections. Digital files of the drilling well locations and production information were obtained through the SAMS database (Lepard et al. 1999).

To display the well locations on a geological map and in three dimensional space, the SAMS well location and database formation information were combined and imported into a three dimensional Microstation 95 format using visual basic programs created by D. Lebel (GSC-Quebec). Each digital well-log file (.las format) was imported into Geographix LogM v. 3-0 (Landmark) and individually interpreted. As well, the location coordinates (UTM NAD83) for each well were input to the same projection as the geological map of Figure 1-3. To create formation isopach maps, ascii files were

WELL SECTION LOCATION INDEX MAP
TRUTCH AND HALFWAY RIVER
 BRITISH COLUMBIA

SCALE
 Kilometres 8 0 8 16 24 Kilomètres

Universal Transverse Mercator Projection
 © Her Majesty the Queen in Right of Canada, 2002
 Projection transversale universelle de Mercator
 © Sa Majesté la Reine du chef du Canada, 2002

Figure 3-1: Location map for the interpreted wells (green dots).

created and combined for each well, including the standardized formation picks plus the well-log location coordinates. The combined ascii file was imported into Zycor ZMap Plus (Landmark) by Phil Lawrence of GSC Calgary.

The well data map coverage (Figure 3-1) is too sparse in areas for reliable computer contouring. To overcome this problem, the formation thickness was plotted at each well-log location and exported as a .dxf file. The .dxf file was imported into Microstation 95 and two 1:100,000 scale plots of the Lower Carboniferous Kiskatinaw and Golata formations were created and hand contoured. The two contour maps were digitized in Autocad and imported to CorelDraw 9.0 for final presentation. The well-log cross-sections of Figure 1-9 were created from GMAPPlus Model Builder v. 3-0. Model Builder incorporated the LogM well-logs and created scaled cross-sections that allow the user to define a datum horizon. To produce the final color cross-sections of Figure 1-9, the sections were imported into Autocad 14 to build the polygons and finally to CorelDraw 9.0 for final presentation.

3.2.2 Well-Log Interpretation

Figure 3-2 is a chart that correlates the regional stratigraphy with the well-log gamma ray and sonic log signatures and a representative synthetic seismogram. The subsurface Triassic formation picks of Dixon (1999) were used to aid in the initial correlations of these subsurface units. As well, the Canadian Stratigraphic Service Ltd. in Calgary Alberta has done detailed well-log analysis and formation picks of selected wells in the Pink Mountain area, for example well D-63-D-94/G2. These CANSTRAT logs add to the reliability of the well-log interpretations for the study area.

From consultation of the above, the most distinct marker horizons on the well-logs are: the Triassic Nordegg Member of the Fernie Formation, the Triassic Doig-Montney Formation contact, the Triassic-Permian unconformity, and the Carboniferous Golata Formation. B. Richards of GSC-Calgary verified the position of the Carboniferous well-log boundaries. The Triassic Nordegg Member top, Baldonnell Formation top, and Charlie Lake Formation base were verified by Gabriella Carrelli of the University of Calgary and J.P. Zonneveld of GSC-Calgary. The preceding evaluations of the entire

Figure 3-2: The subsurface well-log stratigraphy and synthetic seismogram.

well-log sequence ensure consistent well-log picks between this study and other studies that have been done or are in progress.

In the Halfway River area, the only new unit consists of the Cretaceous Cadomin Formation sandstones and conglomerates, which appears as a 100-250 metre thick sequence on cross-sections compiled by Thompson (1989). In his memoir, Stott (1998) documents the presence of the Cadomin Formation in some of the 94B/14 and 94B/15 well-logs. For this study, the Cadomin Formation is included where interpreted on the well-logs across portions of Well Cross-sections 5 and 6 (Figure 1-9). Marking the position of the Cadomin Formation horizon reveals well-log characteristics that distinguish the Monteith from the Gething Formation in the Halfway River area. Projecting the Monteith and Gething formation well-log characteristics northward into the Pink Mountain area would involve a more detailed core analysis study that is beyond the scope of this thesis.

The oldest units on the deepest wells are identified as the Cambrian System quartzite sequences on the CANSTRAT logs (well B-82-F-94G/11). These quartz sandstone and rare carbonate sequences could belong to the quartzite unit that is approximately equivalent in age to the Cambrian Gog Formation (Stott 1991). However, the quartz sandstone sequences are most likely the basal sands of the Dunedin, Stone or Muncho-McConnell formations (well B-10-J-94/G7, Ross et al. 1993). These sandstones are not a typical lithology within the older Devonian formations and the clastic deposition could be the result of structural uplift within the study area during the middle Devonian (Ross et al. 1993).

3.3 Previous Well-Log Research in the Halfway River Area

The extensive well-log study conducted in the Pink Mountain area is an extension of an earlier well-log research paper in the Peace River area by Barclay et al. (1990). Barclay et al.'s (1990) study area covered the Dawson Creek Graben Complex formed over the relict Peace River Arch (Figure 3-3) and consisted of over 1200 wells. During the Late Devonian (?) – Carboniferous in the Peace River area, the deposition of the Stoddart Group was strongly influenced by the formation of steep normal fault structures

Figure 3-3: Carboniferous Dawson Creek Graben Complex and the Stoddart Group isopach map of the Peace River area (modified from Barclay et al. 1990).

above the relict Peace River Arch. Through well-log cross-sections and isopach maps, Barclay et al. (1990) observed the maximum thickness of the Stoddart Group occurring in a rift valley style pattern that matches subsurface maps of the Dawson Creek Graben Complex (Figure 3-3).

In the Pink Mountain area, 150 wells penetrate below the base of the Stoddart Group into older stratigraphy. As seen in Figure 3-1, the data coverage is adequate to indicate thickness change trends within the Stoddart Group of the study area. In Chapter 5, two isopach maps of the Kiskatinaw and Golata formations will be used in conjunction with seismic sections to map the structural configuration of subsurface normal faults within the deeper Devonian-Carboniferous strata.

4. STRUCTURAL GEOLOGY OF THE TRUTCH AND HALFWAY RIVER AREAS

4.1 Generalized Structural Setting of the Cordilleran Deformation

Detailed stratigraphic and structural studies indicate that the present structural style of the Cordillera was governed by three major tectonic events throughout geologic time: 1) Proterozoic to late Triassic extension, 2) Jurassic to Tertiary convergence, and 3) minor Eocene to Oligocene extension. The first stage of extension created an Atlantic-style continental margin, which caused the North American continental plate to stretch and subside resulting in the formation of normal and strike-slip faults in Precambrian rocks throughout the Cordillera (Fermor and Moffat, 1992). These Precambrian basement faults are thought to influence the development of lateral transfer zones or fold culminations during the later Jurassic to Tertiary Columbian and Laramide orogenies that folded and faulted the overlying Paleozoic carbonate and clastic rock sequences during the Pacific plate and North American plate collision (Thompson 1989).

Within the northern Canadian Cordillera, some characteristic features resulting from the mostly compressional deformation are low angle thrust faults that dip west (some are folded), detachment box folds in the hanging wall of the thrust sheet and broad synclines in the footwalls of thrusts (Bally et al. 1966, Dahlstrom 1970). At some thrust sheet boundaries in the southern Canadian Cordillera, lateral transfer zones between folds and faults form, such as the Limestone Mountain Culmination, northwest of Calgary, Alberta (Bégin and Spratt 2002).

4.2 Formation and Configuration of Thrust Fault Systems

During compression, a thrust fault nucleates at a zone of weakness and usually cuts up section along a stair-step or ramp-flat geometry in the direction of tectonic transport. Most commonly, the thrust forms stairs or flats within incompetent strata (shales) bedding planes, and steps or ramps oblique to bedding in competent (sandstones, carbonates) strata (Dahlstrom, 1970). The fault nucleation zone could be a facies change in the strata or a result of the maximum rock folding strain being exceeded (Liu and Dixon 1992, Cobbald 1975).

More important is the presence of older, more steeply dipping faults deeper in the subsurface that are proposed to act as nucleation zones for thrust fault formation (Thompson 1989). For this study, all tectonic models of fault structures will assume the Cordilleran configuration of hinterland dipping duplexes formed from the eastward movement of the Pacific plate against the North American plate during Jurassic to Eocene compressional tectonics.

One of the most common Cordilleran structures is the thrust fault. The volume of rock above it, which is carried by the thrust fault, is termed a thrust sheet. An imbricate fan of thrust sheets form when successive thrust faults branch from the same basal thrust (Fox 1959, Jones 1971; Figure 4-1). Nucleation of new thrust faults occurs when the compressive strain energy required to move the older, more voluminous, steeply dipping or folded thrust sheet is more efficiently transferred to initiating a new, more shallowly dipping fault beneath the older thrust sheet (Boyer and Elliott 1982, Lageson 1984, Cobbold and Gapais 1987). Splaying thrust faults can merge together at a higher stratigraphic level and the rock volume bounded by the enclosing fault surfaces is termed a horse (Figure 4-1).

As observed by numerous studies and first outlined by Dahlstrom (1970; the Cordilleran style rules of deformation), the imbricate fan of thrusts branches upward from a common sole thrust deeper in the subsurface. A duplex system of thrust sheets results when the imbricate fan is bounded by both a basal sole thrust (or floor thrust) and an upper roof thrust. Figure 4-1 outlines the development of a duplex fault system between two glide horizons. The new thrust (dotted line) propagates from the base of the older thrust ramp and eventually ramps up and meets the upper thrust farther east, thus forming the initial horse. This initial horse slides along the new ramp and forms a kink-folded syncline-anticline pair within the horse (Boyer and Elliott 1982, Dahlstrom 1970, Jones 1971).

The thrust fault propagation diagram presented in Figure 4-1 is only one possible model for thrust fault propagation and does not integrate the effects of detachment folding over a basal detachment within the Besa River Formation, which has been proposed by Thompson (1989; Figure 1-5a). The model of Figure 4-1 could partially represent the lower Triassic and Carboniferous stratigraphic horizons farther west of Pink Mountain in the Sikanni area, where the compressional strain was greater during the Laramide Orogeny.

Figure 4-1: Evolution of a Cordilleran thrust fault system. This diagram is derived from Cordilleran examples assuming plane strain and kink folding (**S**-fault slip, **B** - fault angle, modified from Boyer and Elliott 1982).

Interpretation of well-logs immediately north of Two Bit Creek indicates local thickening of the Stoddart Group and Prophet Formation (Figures 1-3 and 1-9). These thick packages could be interpreted as possible duplex structures. If this interpretation were correct, the more resistant Carboniferous stratigraphy would form duplex structures in between the shales of the Doig-Montney formations and the Besa River Formation. In Figure 4-1, the Pink Mountain Anticline would be placed farther east in front of the main duplex structure, where the effects of detachment folding are more dominant. Thus, the more practical model of thrust fault propagation and detachment folding for the Pink Mountain area would be a combination of Figures 4-1 and 1-5a.

4.3 Structural Geology of the Trutch (94G) and Halfway River (94B) Areas

From surface mapping observations, folding is the dominant structural process that occurs within the Triassic and younger strata of the Trutch (94G) and Halfway River (94B) areas, in response to the Laramide Orogeny (Stott 1963; Figures 1-3 and 1-8 – geologic maps). Within the 94G/2 and 94G/3 areas, the folds of Triassic and younger strata in this area are mostly asymmetrical box shaped folds with the steeper limb commonly dipping to the west (Cecile 1999). In areas where the Triassic strata are capped by the thicker, more resistant clastic sequences of the lower Cretaceous, the box folds at this stratigraphic level are broader with more shallowly dipping asymmetrical limbs. In the Triassic cores of these folds, the less resistant shales and carbonates of the Triassic have steeper limbs that are more symmetrical in appearance (Lane et al. 1999), one such example is the Pink Mountain Anticline (Figure 4-2).

Fold plunges are variable within the Pink Mountain – Marion Lake area. Throughout most of the Pink Mountain – Marion Lake area, the surface traces of the fold axes extend for over 24 kilometres and have shallow (eight degrees and less) plunges. These long-axis folds trend approximately 344° - 354° . Locally, however, Cecile (1999) observed abrupt changes in plunge, with steeper (greater than 10°) fold plunges along the Sikanni Chief River (Marion Lake). Also, the fold axial traces change orientation (besides topographic effects) to a more northerly trend in the central areas of the Marion Lake map sheet (Cecile 1999). Once again,

Figure 4-2: View of Pink Mountain looking north. The steeper limb of the Triassic fold is outlined in white (photo by Andrew Miall).

there is no surface evidence for why the fold axes change orientation in some areas. Surface evidence is scant for faulting within the Triassic and younger strata in the southern Trutch area (M. Cecile, pers. comm.). In the Marion Lake map area (94G/3), only four thrust faults with minor displacements are mapped within the Triassic strata. The surface extent of these faults is 2 to 10 kilometres (Figure 1-8 geologic map). The thrusts are related to fault-propagation folds and probably sole within the thicker shales of the Toad-Grayling formations (Doig and Montney formations in the subsurface), or they could sole within the deeper Besa River Formation.

4.4 Subsurface Seismic Resolution in the Trutch area

The quality of seismic data within the Trutch area is highly variable. Figure 4-3 is a location map of all the seismic lines in the study area made available by industry. Poor seismic resolution is partially caused by the processors utilizing incorrect velocity models and under or over-migrating the deeper structures. One of the main causes of poor seismic reflections is the presence of a fractured, bouldered surface in the topographically high areas, which is caused by frost-heaving and slumping of the resistant lithologies of the Triassic to lower Cretaceous stratigraphy (Huber A., and Cecile M. pers. comm.). The rubbly surface attenuates the signal and decreases the strength or completely cancels out reflections from deeper structures.

Another fundamental problem with seismic resolution in the Trutch area is the difficulty in imaging faults within steep limbs of box folds. Also, the combination of faults and steeply dipping bedding planes could reduce the seismic signal to noise ratio to zero (A. Newson, pers. comm.). One such poorly resolved horizon exists from the northern extent of Pink Mountain (Figure 1-7 – north seismic section) to the Sikanni area just west of it (A. Newson, pers. comm.). Multiple seismic surveys shot under a variety of operating parameters have failed to significantly improve the seismic imaging of this difficult area.

The unprocessed seismic lines in Figure 4-3 are colored grey and the three seismic lines used for cross-sections in this study are red, green and orange. A Petro-Canada dip line north of Pink Mountain, a Sigma Explorations Inc. dip line south of Pink Mountain, and a

Figure 4-3: Location map of the digital well and seismic data for the study area.

Petro-Canada strike line along the eastern edge of the central portion of Pink Mountain will be used to determine the deeper structural origin of the study area. To aid in the interpretation, Vlada Avramovic of GX Technologies (formerly of Kelman Technologies) has re-processed the northern and southern seismic sections (Figure 1-7) using pre-stack time and pre-stack depth migration methods to better resolve the deeper stratigraphy and structure. Mr. Avramovic was provided with geological models and well-logs for more accurate velocity determinations and he will include a more detailed summary of the processing workflow in his MSc. thesis (in progress).

Interpretations of the deeper seismic data will follow a general summary of the surface and shallow subsurface geology of Pink Mountain.

4.5 Surface Structural Geology of the Pink Mountain Anticline

Both Pink and Spruce Mountains comprise a box anticline structure with a major axis that appears to be both offset and re-oriented across the Halfway River (Figures 1-1 and 1-3). Older interpretations of Pink Mountain show the box anticline axial trace changing from approximately 344° to 355° at Halfway River (Hage 1944, Taylor 1979). However, new map data and stereonet of the Halfway River area show all fold axes trend between 335° - 340° and plunge direction changes along the Pink Mountain Anticline (Figure 4-4). The fold plunge changes are similar to those documented in the Sikanni River area by Cecile (1999). From both surface and stereonet analysis, the major box anticline that extends northward from Spruce Mountain across Halfway River dies out west of the major Pink Mountain box anticline.

The major box anticline of Pink Mountain is clearly seen at Halfway River and its axis extends northwest with a northwest plunge until its surface expression disappears at Little Pink (Figures 1-8, 4-2 and 4-4). A more detailed stereonet analysis of Pink Mountain is presented in section 4.8. South of Halfway River at Spruce Mountain, the major box anticline axis is accompanied by five minor fold axes that appear to die out 1.5 kilometres to the southeast whereas the major axis continues southward and plunges southeast beneath the surface at 56° $57'$ north latitude (Figure 1-3).

Figure 4-4: Map of Pink and Spruce Mountains at Halfway River showing stereoplots in the three main fold domains. Backthrusts represent the western boundaries of Domains 1 and 3, dashed grey line marks the boundary of Domain 2.

The folding style of the Pink Mountain Anticline varies northward from an asymmetric box anticline at Halfway River (Sections 2 and 3, Figure 1-3) to a broader anticline with shallower limb dips at Little Pink (Sections 5 to 7, Figure 1-3). The fold structure of Spruce Mountain is mostly an asymmetric box anticline with a steeper west limb and a mostly planar, shallowly dipping eastern limb (Section 1, Figure 1-3).

Surrounding Pink Mountain, vegetation cover on the Buckinghorse Formation shales obscures possible faults (Figure 1-3). Only two minor faults, an eastern thrust and a backthrust have been observed along Moose Lick Creek, 4 km northeast of Pink Mountain. Both faults are within a gully cut into the Buckinghorse Formation and have less than 400 metres of displacement. If this displacement were greater than 400 metres, older rocks would crop out. The structural origins of the Pink Mountain and Spruce Mountain anticlines cannot be deduced from surface observations, so two cross-sections across the northern and southern ends of the Pink Mountain Anticline were compiled based on the highest confidence surface mapping, well-logs, and seismic data (Figures 1-7 and 1-8).

4.6 Subsurface Structural Interpretation of Pink Mountain

The surface expression of the Pink Mountain Anticline has resulted in the drilling of nine exploration wells along Pink Mountain during the late 1960's to 1990's. Interpretation of four well-logs drilled on the western limb of the Pink Mountain Anticline has revealed a repetition of the Carboniferous Golata Formation to the Triassic Charlie Lake Formation packages (Figure 4-5). This repeat reveals the occurrence of a fault beneath Pink Mountain that was not detected by extensive surface mapping. In conjunction with nearby seismic interpretations, the fault is interpreted to be a backthrust (termed the Two Bit Creek Fault) obscured by talus along the western limb of the Pink Mountain Anticline (Figure 1-3, Sections 1 to 11).

The well-log interpretation of Pink Mountain initiated a completely different surface structural interpretation of the Pink Mountain Anticline and its relationship to the southern Spruce Mountain Anticline. Section 3 of Figure 1-3 reveals the presence of a second fault west of the Two Bit Creek backthrust. Both wells C-71-D and D-63-D 94G/2 of Section 3

Figure 4-5: Gamma Ray well-log interpretation of two selected wells at Pink Mountain.

were corrected for bottom hole position and deviation by consulting Mr. Ron Stefik of the British Columbia Oil and Gas Commission. Balancing of Sections 1 to 7 and re-analysis of the surface geology and airphotos of Pink Mountain have led to the interpretation of an en échelon pair of backthrusts in the Halfway River area of Pink Mountain (Figure 1-3).

4.7 Surface Identification of the Pink Mountain Backthrust Faults

On the western flanks of both Pink and Spruce Mountains, surface identification of a backthrust was limited to two traverse areas on western Pink Mountain that exhibited a high degree of fracturing of the Monteith-Gething formations. This fracturing may have been related to brittle deformation of the resistant sandstones of the Monteith-Gething formations during the folding stage of the Pink Mountain Anticline. At Halfway River, the surface expression of Two Bit Creek Fault is obscured by talus and tree growth. The only evidence for a possible fault in this area is the 1.2 km² rock slump and the steep-limbed minor folds in the road exposures on Pink Mountain east of the slump (Figure 1-3).

Confirmation of the surface position of the Two Bit Creek backthrust and where it crosses Pink Mountain came from 1944 air photographs taken before the construction of the communications tower (Figure 4-6). In these old airphotos, a lineament is seen trending southeast across Pink Mountain from the position of the (later) wells that intersect the repeated section (inferred backthrust) and extends to the eastern area of the rock slump and steep-limbed minor folds. The construction of the road leading to the communications tower at the summit of Pink Mountain during the 1950's obscured the surface expression of the Two Bit Creek Fault across Pink Mountain near the Halfway River (Figures 1-3 and 4-6).

4.8 Stereonet Analysis of Pink and Spruce Mountains

The spatial variation in fold axis orientation across the study area was analyzed using Pangaea Scientific's SpheriStat 2.1c software. The area was divided into domains and sub-domains, then strikes and dips of bedding were used to calculate the mean fold axis orientation for each sub-domain. Stereonets for the sub-domains in the vicinity of Halfway River are shown in Figure 4-4.

Figure 4-6: Composite airphoto mosaic of Pink Mountain from photos taken in 1944. The Two Bit Creek Fault can be seen in the dashed outline (modified from Department of Energy Mines and Resources Canada).

A major domain boundary between the Pink and Spruce Mountain structures is the Two Bit Creek Fault that cuts across the southern end of Pink Mountain. As seen in Figure 4-4, two major and one minor fold domains exist in the Halfway River area: two domains with northwest trends, the main Pink and Spruce Mountain structures (Domains 1 and 3), and a much smaller domain (Domain 2) with a more northerly trend, located 1 to 2 kilometres south of the summit of Pink Mountain (Figure 4-4).

With mapping of the Two Bit Creek Fault across Pink Mountain, it becomes clear that the Pink Mountain Structure (Domain 1) and the Spruce Mountain Structure (Domain 3) are two separate, en échelon folds (Figure 4-4), not a single anticline with an offset axial trace. In the vicinity of Halfway River, the Pink Mountain Structure plunges south-southeast, and the Spruce Mountain Structure plunges north-northwest, dying out just west of the summit of Pink Mountain.

4.9 Deep Structural Interpretation of the Pink Mountain Area

The final re-processing (pre-stack depth migration) of the northern and southern seismic sections resulted in improved resolution down to depths of 5 kilometres (Figure 1-7). Nearby wells were projected into the seismic sections and the tops of units were correlated laterally across the section. Seismic character was used to correlate events across faults and zones with poor resolution. The oldest identifiable unit, the Devonian Stone Formation, is imaged on the northern seismic section. On the southern section, the resolution fades below the Dunedin Formation and only a tentative identification of the Cambrian to Devonian reflector can be made because this interval has been interpreted to thin approximately 40-50 kilometres northeast of Pink Mountain (M. McMechan pers. comm.).

Interpretations of both seismic sections reveal that the Pink and Spruce Mountain structures occur between an eastern thrust fault and a western backthrust. Both faults have less than 1 kilometre of displacement (Figures 1-7 and 1-8). On both sections, the east-dipping backthrust soles into the main thrust fault. The Pink and Spruce Mountain anticlines are contained within this large scale pop-up structure, east of the Rocky Mountain Foothills, that extends from Cypress Creek to north of the Sikanni Chief River (Figures 1-3 and 1-8).

Beneath and to the east of the pop-up structure, deeper west-dipping normal faults are interpreted on both seismic sections. These steeper faults interrupt the Cambrian-Devonian events and appear to extend up to the Jurassic-Cretaceous Monteith-Gething level (Figure 1-7). On both seismic sections, above the normal faults, there is a westerly increase in thickness of the Carboniferous Stoddart Group, the Triassic Upper Montney Formation, and possibly the Monteith-Gething formations. This suggests that several episodes of extensional tectonism and subsidence alternated with periods of quiescence before the onset of compression. During the Laramide Orogeny, the normal fault east of the pop-up structure was mildly re-activated as a thrust fault (Figures 1-7 and 1-8).

4.10 Balanced Cross-sections of the Pink and Spruce Mountain Anticlines

The northern and southern seismic sections have been depth-migrated (by Vlada Avramovic) and tied to the nearby wells, so it was possible to test the seismic interpretations using bed length and area balancing techniques. The line interpretations of Figure 1-7 were digitized and converted to Microstation 95 format in a Universal Transverse Mercator (NAD83) projection. Both seismic interpretations were bed-length balanced to within 1.5% error and area balanced to within 4.0% error in Microstation 95 (Figure 1-8). Minor adjustments were made to the line work in Microstation 95, but these changes had a negligible effect on the seismic interpretations shown in Figure 1-7.

The 13.5 kilometre long northern seismic section exhibits 1.3 kilometres of shortening of strata above the Besa River Formation. This is equivalent to a shortening percent ($[(l_o - l_f) / l_o] \times 100$) of 8.8%. However, the strata below the Besa River detachment experienced less than 200 metres of net shortening (<1.5%; Figure 1-8). This agrees with previous proposals that, in the eastern part of the Rocky Mountain Foothills, the Besa River Formation shales absorb most of the tectonic stresses and the older strata are relatively undeformed (Thompson 1989).

The 32 kilometre long southern seismic section exhibits 1.0 kilometre of shortening (3.0%) of strata above the Besa River Formation (Figure 1-8). Strata below the Besa River Formation experienced less than 1 kilometre of shortening by folding, but they record a net

extension of approximately 4 kilometres (14.3%) due to the significant normal faults in the deep part of the section. The imbalance in shortening between the northern and southern sections would likely be remedied if both sections were extended farther to the west, where duplex structures and folding account for more shortening of units beneath the Besa River detachment (D. Spratt, pers. comm.).

5. TECTONIC HISTORY OF THE TRUTCH AND HALFWAY RIVER AREAS

5.1 Pink Mountain Area in Relation to the Peace River Arch

The present day structural configuration of the Pink Mountain Anticline could be the result of a combination of tectonic events related to the origin and development of the Peace River Arch south of the study area. The exact origins of the Arch are unclear and are still a matter of debate among scientists. Two popular theories on the origin of the Peace River Arch in the Precambrian are: a failed rift zone and a continental fracture system adjacent to a spreading oceanic arc (O'Connell et al. 1990). The failed rift scenario has several complications including the lack of thermal decay (causing subsidence) of the topographically high Arch from the Precambrian to Devonian. Physical evidence for a fracture system originating from a possible nearby oceanic spreading margin is the offset of the east-west trending Peace River Arch across a northwest-southeast feature (Figure 5-1a). The offset in the Peace River trend resembles transverse faulting across spreading ridges associated with modern oceanic spreading margins (O'Connell et al. 1990, Wilson and Williams 1979). Direct evidence for the oceanic spreading margin is not presented in the research literature.

5.2 Precambrian to Cambrian Tectonic History

The Precambrian basement in the study area is composed of five main domains from southeast to northwest, the Buffalo Head, Chinchaga, Ksituan, Slave, and Hottah, (Figure 5-2) with boundaries that appear to have been re-activated throughout the Phanerozoic (Ross et al. 1993). Throughout the Upper Proterozoic to Middle Cambrian, the Peace River Arch was a positive tectonic feature that provided most of the source material for the Precambrian Windermere Supergroup and Cambrian Gog Group (O'Connell et al. 1990, McMechan 1990). North of the Peace River Arch, the Gog Group was deposited on an uneven (block faulted?) Precambrian basement surface in an extensional tectonic environment and hence the thickness of the formation varies laterally (Puch 1975, O'Connell et al. 1990, McMechan 1990). Deeper east-west private industry seismic sections reveal thick packages of laterally discontinuous reflectors that resemble basin fill sediments at approximately 3 to 6 kilometres depth.

Figure 5-1: a) Structure contour map and b) cross-section of the Peace River Arch Precambrian surface (modified from Trotter 1989).

Figure 5-2: Basement domains of western Alberta and northeastern British Columbia (modified from Ross et al. 1993)

5.3 Ordovician and Silurian Quiescence

During the Ordovician and Silurian periods, the Peace River region was mostly a passive continental margin of the North American Craton. The majority of the sediments were carbonates deposited on an extensive tropical platform surrounding the Peace River Arch. These carbonates were periodically interrupted by silica-rich sediment detritus shed from the Peace River Arch and other topographic highs during brief episodes of tectonic activity. The mostly carbonate sediments of the Lower Ordovician Kechika Group (Figure 2-2) were deposited on a gentle continental slope and broad platform that covered a large area surrounding the study area (O'Connell et al. 1990, Norford 1990).

At the beginning of the Middle Ordovician a transition area developed, between the shallower MacDonald Platform in the east and the deeper Ospika Embayment basin environments in the west (Figure 2-3), that accounts for the east-west lithological variations in the study area (Cecile and Norford, 1979). From the Middle Ordovician to Lower Devonian, the shale-dominant Road River Group was deposited within the Ospika Embayment while the MacDonald Platform hosted the mostly carbonate units of the Skoki, Beaverfoot, Nonda and Muncho-McConnell formations (Figures 2-2 and 2-3, Cecile et al. 1997, Norford 1990). Periodic minor tectonic uplift episodes throughout the Middle Ordovician to Late Silurian were responsible for the lateral movement of the platform-shelf facies transition and the extensive regional unconformities in the units between the Skoki and Muncho-McConnell formations (Cecile and Norford 1979).

5.4 Devonian and Carboniferous Tectonism and the Dawson Creek Graben Complex

At the beginning of the Middle Devonian, a period of tectonic uplift in the Peace River area could have been the cause of the widespread deposition of the Granite Wash lithosome (Trotter 1989). The thickness of the Granite Wash can reach 100 metres within graben structures near the crest of the Peace River Arch and on broad depositional planes north of the Peace River Arch (O'Connell et al. 1990). During the Late Devonian the depositional environment northwest of the Peace River Arch was predominantly a platform setting, which resulted in the deposition of the mostly carbonate facies of the

Dunedin and Stone formations (Figure 2-2, Taylor and MacKenzie 1970). This continual sediment deposition resulted in the burial of the Peace River Arch by the Lower Carboniferous (Figure 5-1b, O'Connell et al. 1990). Northwest of the Peace River Arch, a higher rate of subsidence in the Middle to Upper Carboniferous resulted in the deposition of thicker deep-water units such as the Besa River and Prophet formations (Figure 2-2, Bamber and Mamet 1978). The thickness variations within the Prophet Formation equivalents (Banff and Pekisko formations) could have been controlled by the re-activation of the fracture system along the northern portion of the Peace River Arch (Figure 5-1a, O'Connell et al. 1990). During the deposition of the Prophet Formation, a prominent east-west elongated basin termed the Peace River Embayment was formed in the area above the buried Peace River Arch (Figure 5-3, Barclay et al. 1990).

Throughout the Carboniferous, the Peace River Embayment experienced tectonic subsidence along some of the old fracture systems of the Peace River Arch (Figure 5-3). Subsidence was greatest within the Peace River Embayment relative to the rest of the Western Canada Sedimentary Basin. Approximately 185 km southeast of Pink Mountain, an extensive fault system termed the Dawson Creek Graben Complex affected the Devonian-Carboniferous and older strata during the deposition of the Stoddart Group and Belloy Formation (Figure 5-4a).

5.4.1 Thickness Variations within the Stoddart Group in the Peace River Area

The Dawson Creek Graben Complex contains the extensive northeast-southwest trending Fort St. John Graben and well logs across this system document changes in elevations of the tops of the upper Prophet (or Debolt) Formation (Barclay et al. 1990). Also, abrupt thickness variations are observed within the Kiskatinaw and Golata formations of the Stoddart Group (Figure 5-4b). In the Peace River area, the Stoddart Group has a vertical thickness variation of 100 to 200 metres over a horizontal distance of two kilometers (Barclay et al. 1990). During the Carboniferous, sediment deposition was limited to an embayment spanning from west central Alberta to northeastern British Columbia, indicating a phase of relatively local extension (O'Connell et al. 1990, Barclay et al. 1990). Figure 5-4b is a schematic portrayal of the deposition of the Stoddart Group

Figure 5-3: Carboniferous-Permian geological and tectonic elements in northeastern British Columbia and Alberta (modified from Barclay et al. 1990).

Figure 5-4: Schematic diagram of a) the Carboniferous-Permian Dawson Creek Graben Complex and b) the depositional history of the Stoddart Group (modified from Barclay et al. 1990; see Figure 2-2 for stratigraphy).

formations across these proposed graben structures. The most important factor of Figure 5-4b is the absence of the Taylor Flat carbonate facies surrounding the Fort St. John Graben. Figure 5-4b indicates that the Taylor Flats Formation is restricted to the immediate graben areas, which explains why this formation is mostly absent from the subsurface stratigraphy of the Trutch area (B. Richards, pers. comm.).

5.4.2 Stoddart Group Depositional Style in the Pink Mountain Area

Figure 5-5 shows two isopach maps constructed for the Carboniferous Golata and Kiskatinaw formations of the Stoddart Group in the Trutch-Halfway River map areas. The first observation that is apparent is the similarity in the pattern of thickness variations of the Stoddart Group lithologies compared to the isopach map of Barclay et al. (1990) (Figure 3-3). The most noticeable feature of the Figure 5-5 maps is the 600+ metre thickness of the combined Kiskatinaw and Golata formations in the south. There is an abrupt increase in the thickness of the Stoddart Group across an east-west trending zone in the southern portion of the maps. In the northern parts of the Figure 5-5 maps, the thicker portions of the Stoddart Group form elongate lobes or valleys (marked by thick dashed lines) that trend approximately east-west. These valleys are separated by thin Stoddart Group horizons. One explanation for these thickening trends is satellite graben structures off the main southern graben (Figure 5-5a).

The nine well cross-sections of Figure 1-9 also show the thickness change trends of the Carboniferous Stoddart Group seen in the isopach maps. The only areas open to interpretation on the east-west well cross-sections are the extreme western ends, which could have structural repetition (marked on sections) of the Stoddart Group (Sections 1 - 6 Figure 1-9). Sections 8 and 9 of Figure 1-9, along with the two isopach maps, indicate two possible minor east-west grabens controlling the deposition of the Stoddart Group (Figure 5-5 and Figure 1-9). The zero edge of deposition on both isopach maps could indicate areas of subsided normal fault blocks in the west and south.

Figure 5-5: Isopach maps (contours in metres) of the Carboniferous a) Kiskatinaw and b) Golata formations in the Trutch and Halfway map areas. Seismic lines shown in orange; dashed grey lines are possible subsurface faults.

5.4.3 Evidence for Major and Minor Subsurface Normal Faults at Pink Mountain

Concrete evidence for the influence of subsurface normal faults affecting the thickness of the Stoddart Group can be seen in Figures 1-7 and 1-8. The seismic interpretation of Figure 1-7 for lines north and south of Pink Mountain clearly shows the westward thickening of the Stoddart Group above a deeper subsurface normal fault, which down-drops the Devonian-Carboniferous and older strata along the western portion of both seismic sections. From inferred positions on the northern and southern seismic lines, the overall trend of this subsurface normal fault east of Pink Mountain (termed the Pink Mountain Fault) is 340° , which approximately follows the strike of the beds on Pink Mountain, north of the Halfway River and the en échelon backthrusts (Figures 1-3 and 1-7).

The subsurface configuration of the Stoddart Group outlines the complex depositional environment of subsidence and erosional uplift of the graben system during the later Carboniferous and early Permian (Figure 5-3 and 5-4b). Additional steep subsurface faults may exist farther west, beneath the Pink Mountain Anticline (similar to the eastern “satellite” grabens of Figures 5-3 and 5-4a) and influence the structural geology of the younger units at Pink Mountain. Proof of the existence of one such minor fault in the Halfway River area can be seen in an along-strike seismic section donated by Petro-Canada just east of Pink Mountain (Figure 5-6).

From the seismic interpretation, the Two Bit Creek backthrust beneath Pink Mountain soles into the Besa River Formation and appears to be affected by the presence of a steeper fault that displaces Devonian and older strata. This steeper fault is termed the Elbow Creek Fault because of its close proximity to Elbow Creek (Figure 1-3). The position of the deep fault coincides with the area of en échelon backthrusts in the younger strata above it (Figures 1-3 and 5-6). The Devonian and younger strata form a broad anticline above the Two Bit Creek backthrust south of the Elbow Creek Fault. The strike of the Elbow Creek Fault cannot be constrained by a single seismic section, but it may be west to northwest trending. Similar faults may also exist in Devonian and older strata to the north, accounting for the east-west satellite grabens interpreted on the Stoddart Group

Figure 5-6: North-South seismic section showing the Two Bit Creek Backthrust and the deeper Elbow Creek Fault (seismic section donated by Petro-Canada).

isopach maps of Figure 5-5.

5.4.4 Late Carboniferous to Permian Paleosurface

The Late Carboniferous to Permian subsurface stratigraphy in the Pink Mountain area contains a lateral discontinuity between the eastern Prophet Formation carbonates and the Stoddart Group sandstones and shales that occupy the southern and western parts of the area (Figure 5-5). This discontinuity coincides with the Pink Mountain Fault, which intersects another east-west normal fault near Cypress Creek in the southern part of Figure 5-5 in the eastern Halfway River map area. Both the Pink Mountain and another northwesterly trending fault along Cameron River (termed the Cameron River Fault) are clearly seen on the southern seismic section and balanced cross-section of Figures 1-7 and 1-8.

The Carboniferous “High” east of the zero edge of the Stoddart Group in Figure 5-5 can range in height from 50 to 400 metres and is clearly seen in the interpreted southern seismic section of Figure 1-7 and balanced southern cross-section of Figure 1-8. It is constrained by the northwest-southeast and east-west trending major faults and may represent a resistant carbonate step or buttress that focused tectonic stresses in the overlying rocks.

5.5 Uncertain Tectonism during the Permian to Triassic

The duration of graben movement throughout the Late Carboniferous-Permian is unclear. Thickness variations on a minor scale exist throughout the Permian Belloy Formation, but with small differences in elevations of tops between wells drilled within the Dawson Creek Graben Complex (Barclay et al. 1990). This indicates a reduction in the amount of fault activity and most of the Triassic strata were deposited on a relatively passive continental margin (Gibson and Edwards 1990, O’Connell et al. 1990). However, thickness variations exist within the Triassic Upper Montney Formation that are similar to the isopach patterns of the Carboniferous Stoddart Group within the Pink Mountain area (Figure 5-7).

As seen in Figure 5-7b, the Triassic Lower Montney Formation exhibits a mostly

Figure 5-7: Isopach maps (contours in metres) of the a) Upper Triassic Lower Montney and b) Lower Montney formations in the Trutch and Halfway map areas.

sheet-like depositional pattern with minor east-west channel sequences that suggest a lessening of the graben activity compared to the Carboniferous (Figure 5-5). The depositional pattern of the younger Upper Montney Formation (Figure 5-7a) suggests periodic rejuvenation of tectonic activity along the older fault systems during the Early Triassic. In Figures 1-7 and 1-8, the Montney Formation thickens west of the Pink Mountain Fault in both the northern and southern seismic sections. Note that it is only the Upper Montney Formation that thickens in the northern seismic line (Figures 1-7 and 1-8). The Montney Formation could not be subdivided in the southern seismic section.

Periodic minor rejuvenation of the subsurface normal faults must have persisted as late as the Lower Cretaceous, due to the thickening of the Monteith-Gething formations west of the Pink Mountain and Cameron River Faults. The thickness change of the Monteith-Gething formations across the Cameron River Fault is the easiest to detect on the southern seismic line. Stratigraphic interruptions of the Carboniferous to Cretaceous units occur across both faults. The Pink Mountain Fault shows clear thrust offset at the Monteith-Gething level, indicating that this original normal fault has been partially inverted under compression but the Monteith-Gething thickness change across the Cameron River Fault rules out purely compressional re-activation (Figures 1-7 and 1-8).

5.6 Jurassic to Tertiary Compressional Tectonics

5.6.1 Deposition of the Fernie and Minnes Group and the Columbian Orogeny

The passive continental margin platform environment continued into the Early Jurassic with the deposition of the Lower Fernie Group in the study area. The Lower Fernie Group (Nordegg Member) consists of calcareous shales and limestones, which are interrupted by a regional erosional unconformity related to the Columbian Orogeny (Poulton et al. 1990). The non-calcareous shale-dominated Upper Fernie Group was deposited in a basin formed east of the accreted exotic terranes of the Canadian Cordillera. This elongate eastern foredeep basin resulted from the uplift and thrust faulting during accretion of the exotic terranes (Stott 1998).

In the Middle to Late Jurassic, the seaway between the North American Craton and the Canadian Cordillera narrowed and became shallower in response to the compressional tectonics of the Columbian Orogeny (Stott 1998). This resulted in the deposition of the Jurassic-Cretaceous Minnes Group strata (Figure 2-2). The dominantly clastic Minnes and Bullhead groups are sourced from both the North American Craton and the Canadian Cordillera (Stott 1998). In the Pink Mountain area, the Buckinghorse to Dunvegan formations (Fort St. John Group) were deposited during the waning stages of the Columbian Orogeny (Stott 1991).

5.6.2 The Laramide Orogeny and the Timing of Faults at Pink Mountain

The Laramide Orogeny initiated during the Cenomanian stage of the Cretaceous and compressional tectonics began the folding of the Carboniferous to lower Cretaceous strata above the Devonian-Carboniferous Besa River Formation. This 1.5 kilometre thick sequence of shales, four kilometers deep in the subsurface, acts as a detachment zone for folding and a décollement zone for Laramide thrust faults that cut upsection into the younger stratigraphy (Thompson 1989).

In the Pink Mountain area, continued compression throughout the Upper Cretaceous resulted in the formation of the Lily Lake Thrust, which ramps to the surface in the vicinity of Lily Lake, east of Pink Mountain (Figure 1-3). The Lily Lake Thrust is an eastern splay of a subsurface thrust fault located within the Marion Lake map area (94G/3). A subsurface thrust fault is inferred to exist in the eastern half of the Marion Lake map area based on interpretation of industry seismic data. On both the northern and southern seismic lines, the Lily Lake Thrust extends eastward along a décollement zone within the Besa River Formation beneath Pink Mountain and forms a ramp through the Triassic strata near Lily Lake (Figures 1-3, 1-7 and 1-8). The position and configuration of the Lily Lake Thrust appears to be completely controlled by the presence of the deeper Carboniferous Pink Mountain Fault.

The Pink Mountain Fault has west-side-down offset of less than 500 metres, and this step may have acted as a buttress or stress concentrator, causing the Lily Lake Thrust

to cut upsection from the Besa River décollement, through the Triassic and younger strata just east of Pink Mountain.

5.6.3 Changing Style of the Lily Lake Thrust

Palinspastic restoration of the southern seismic section portrays the Lily Lake Thrust as forming above and to the west of the Pink Mountain Fault, and thus, the two faults are clearly separate entities (Figure 1-8). However, the northern palinspastic restoration shows that the Lily Lake Thrust and the Pink Mountain Faults are in very close proximity to each other as their strikes converge northwards (Figure 1-8). The possibility exists that the lower portion of the Lily Lake Thrust ramp was initially the upward continuation of the Pink Mountain Fault. Although the thickness variation across the fault at the Stoddart level is not as abrupt as in the southern section, the eastern limit of the thickening occurs near this fault. Detailed sequence stratigraphic work would need to be done to determine how much (if any) normal movement occurred on the Lily Lake fault before compression began.

The folds are notably tighter and the Lily Lake Thrust and backthrust trajectories are steeper in the northern section. There they have developed right above the Pink Mountain normal fault, where the stress concentration associated with it would have been greater.

5.6.4 Relationship between Cretaceous and Carboniferous Faults

During the Laramide Orogeny, thrusting and detachment folding were the dominant structural responses to the compressional stresses in the Trutch and Halfway River areas. Steeper than average cutoff angles in the Mesozoic strata are the only evidence that detachment folding of these units may have begun before ramping of the Lily Lake Thrust.

The Lily Lake Thrust and the Two Bit Creek and Spruce Mountain Backthrusts formed a 30+ kilometre long pop-up structure above the “step” in the Carboniferous stratigraphy produced by displacement on the Pink Mountain normal fault (Figures 1-7 and 1-8). The Carboniferous buttress would have impeded translation of the Lily Lake

Thrust sheet toward the foreland, increasing the probability that backthrusts and west-verging folds would develop as compression continued.

5.6.5 Origin of the Two Bit Creek and Spruce Mountain en échelon Backthrusts

The Two Bit Creek Backthrust has its maximum displacement of approximately one kilometre in sections 2 to 5 of Figure 1-3. Its displacement decreases to less than 200 metres in the northern cross-section (Figure 1-8), and it also dies out to the south, between section 1 (Figure 1-3) and the southern cross-section (Figure 1-8). The Spruce Mountain Backthrust has its maximum displacement of about 400 metres just north of section 1 and it decreases to 300 metres in section 2, 200 metres in section 3, and dies out between sections 3 and 4 (Figure 1-3). The Spruce Mountain Backthrust also dies out southward from section 1, as it has less than 100 metres of displacement in the southern cross-section (Figure 1-8). Both faults probably nucleated near the position of maximum displacement and propagated laterally, with the Spruce Mountain Backthrust dying out northward west of where the Two Bit Creek Backthrust has its maximum displacement. Similarly, The Two Bit Creek Backthrust dies out southward just east of where the Spruce Mountain Backthrust has its maximum displacement. They are excellent examples of en échelon faults, but en échelon backthrusts are uncommon in northeastern British Columbia, so why did they form at Pink Mountain?

The cause of the localization of the backthrusts near the Halfway River appears to be the Elbow Creek Fault, a steep transverse fault, which is clearly seen in Figure 5-6 and section 8 of Figure 1-3. The Elbow Creek Fault intersects and is nearly perpendicular to the Pink Mountain Fault, dividing the area into blocks. During Laramide compression, the Spruce Mountain Backthrust nucleated on the southern block, and the Two Bit Creek Backthrust nucleated on the northern block

The Pink Mountain Anticline may have initially formed at depth as a low amplitude, doubly plunging, symmetrical anticline with little to no deflection of its fold axis. As compression continued, the folds tightened and backthrusts nucleated. During the later stages of the Laramide Orogeny, the Two Bit Backthrust propagated southward across the Pink Mountain Anticline and died out just southeast of Pink Mountain at

Halfway River. As a result, the Pink Mountain Anticline became divided into separate structures in the hanging wall and footwall of the Two Bit Backthrust. At the same time, the Spruce Mountain Backthrust propagated north of the Halfway River and died out just north of Section 3 in Figure 1-3.

5.6.6 Cenomanian to Present Configuration of Pink Mountain

During the waning stages of the Laramide Orogeny, the en échelon backthrusts divided the Pink Mountain Anticline into two separate doubly plunging structures, termed the Pink Mountain Anticline north of the Halfway River and the Spruce Mountain Anticline south of the Halfway River (Figure 1-3). As well, both structures developed steeper western limbs due to movement on the en échelon backthrusts. The en échelon backthrusts explain why the fold axes do not line up along strike (Figure 4-4).

5.7 Generalized Synopsis of the Pink Mountain Tectonic History

Figures 5-8 and 5-9 show an overall generalized explanation of the Pink Mountain tectonic history, from the Carboniferous to present. These figures stress the importance of the Carboniferous faults and their influence on the younger sediment deposition and structural geology. In general, the Pink Mountain area experienced extensional tectonics during the Carboniferous, that down-dropped a portion of the paleosurface to the south and west, which was later infilled by the Stoddart Group (Figure 5-8a, b). This tectonism could be the northernmost extent of the Carboniferous Dawson Creek Graben Complex recognized in the Peace River Area (Barclay et al. 1990). The extensional tectonics at Pink Mountain were intermittent throughout the deposition of the Carboniferous Stoddart Group to lower Cretaceous Monteith-Gething formations (Figure 5-8c, d, Figure 5-9e). The resultant step in the lower Cretaceous paleosurface would have coincided with the Pink Mountain Fault (and possibly the Cameron River Fault). The Elbow Creek Fault and other transverse faults may not have extended up to the surface at this stage.

During the Laramide Orogeny, the Triassic to Cretaceous stratigraphy was detachment folded over the Besa River Formation and the Lily Lake Thrust began to ramp up through younger strata. The Pink Mountain Anticline may have initiated as a

Figure 5-8: Tectonic history of the Pink Mountain area; Carboniferous to Lower Cretaceous (Aptian).

Figure 5-9: Tectonic history of the Pink Mountain area; Cretaceous (Albian) to present.

single doubly plunging anticline (Figure 5-9f). Continued compression during the Tertiary caused the initial Two Bit Creek and Spruce Mountain backthrusts to nucleate on either side of the subsurface Elbow Creek transverse fault (Figure 5-9g) as part of a larger scale pop-up structure bounded in the east by the Lily Lake Thrust. The two backthrusts propagated laterally and the Two Bit Creek Fault cut Pink Mountain at a slightly oblique angle, dividing it into the separate Pink Mountain and Spruce Mountain Anticlines. Continued erosion to the present removes the younger Monteith-Gething strata along rivers and gullies to expose the older Triassic strata in the cores of both Pink Mountain and Spruce Mountain Anticlines (Figure 5-9 H). This erosion plus Quaternary to recent cover obscures evidence for the two backthrusts and a possible surface expression of the Pink Mountain Fault.

CHAPTER 6. CONCLUSIONS

Pink Mountain is an anomalous structure that, despite its simple “rounded hulk” surface appearance, raises many questions upon closer examination about the age of the exposed lower Cretaceous stratigraphy, the tectonic origin of the mountain, and how the structural style changes along strike. The first problem the geologist encounters is determining the exact age and formation nomenclature of the sandstones and assorted conglomerates that cap Pink Mountain. Several paleontologists have tried and failed to find preserved fossils at or near the base of these sandstones and have included all of these sequences within the Lower Cretaceous Gething Formation (Gibson 1992, Stott 1998). Dave McNeil of GSC-Calgary has identified *Glomerspirella* species, in samples from the base of these sandstones, which are Upper Jurassic to Lower Cretaceous in age. This confirms the existence of the Upper Jurassic Upper Fernie Formation and the Upper Jurassic to Lower Cretaceous Monteith Formation at the base of these sandstones at Pink Mountain.

However, the problem of distinguishing the Monteith and Gething formations from each other remains because of their similar depositional environments and the fact that the intervening conglomerate horizon associated with the Cadomin Formation cannot be traced laterally across the area. At Pink Mountain, the Gething Formation lies unconformably on top of the partially (or completely?) eroded Monteith Formation. Until evidence is found to separate these two formations, the stratigraphy that caps Pink Mountain is assigned to the undivided Monteith-Gething formations.

One of the most important discoveries of this study is that Pink Mountain, previously interpreted by Hage (1944), McLearn (1950), and Taylor (1979) to be a single anticline, actually consists of two separate box anticlines separated by an en échelon backthrust in the vicinity of Halfway River (Figure 1-3). These anticlines are hereby termed the Pink Mountain Anticline (north of Halfway River) and the Spruce Mountain Anticline (mostly south of Halfway River). The Pink Mountain area has a complex tectonic history involving episodes of Carboniferous to Triassic and possibly Lower

Cretaceous normal faulting that alternated with periods of quiescence prior to compression during the Laramide Orogeny (Late Cretaceous to Paleocene).

Isopach maps of the Carboniferous Stoddart Group indicate the presence of normal faults, within the Carboniferous and older strata, that have been formed by extensional tectonics and may be related to the formation of the Dawson Creek Graben Complex in the Peace River area (Figures 5-4 and 5-5) because they have a similar geometry and timing. Portions of the Halfway River, Elbow Creek, Cameron River, and Sikanni Chief River all follow along the trends of the major normal faults identified in seismic sections and isopach maps. Further study of the subsurface geology beneath these major rivers should provide insight into the extent of these and other faults beyond the study area.

The sub-Permian paleosurface in the Trutch-Halfway River area is a lateral disconformity above the Prophet Formation, which was deposited on the Beaton High area to the east, and the Stoddart Group, which was deposited on the down-dropped normal fault blocks to the west (Figures 5-8a and 5-8b). Based on interpretation of the southern seismic section, several phases of extensional tectonic activity appear to have occurred prior to Laramide compression. Extension affected deposition of the Carboniferous Stoddart Group, the Triassic Upper Montney Formation, and possibly the Lower Cretaceous Monteith-Gething formations (Figure 1-8).

During the Laramide Orogeny, detachment folds developed above the Devonian-Carboniferous Besa River Formation detachment horizon. Folds and thrust ramps nucleated above and to the west of deeper subsurface normal faults within the Carboniferous and older strata. The Pink Mountain Fault, a major northwest trending west-side-down normal fault, created a step or buttress that impeded motion on the Besa River detachment and caused the Lily Lake Thrust to ramp up-section over it. West of this step, a pair of backthrusts nucleated on either side of a minor east-west trending Carboniferous fault and propagated across it in an en échelon pattern. The Two Bit Creek Backthrust ramped laterally across the area and separated the Pink Mountain and Spruce Mountain structures (Figures 4-4 and 5-9h).

Based on seismic interpretations, the Pink and Spruce Mountain structures occur within a thrust-backthrust pop-up structure that extends from Robertson Creek to the Sikanni Chief River (Figure 1-7). The faults of this pop-up structure sole into the Devonian-Carboniferous Besa River Formation and both the northern and southern seismic sections confirm Thompson's (1989) observation that the Besa River Formation acts as a thin-skinned detachment horizon. However, the development and localization of this pop-up is directly related to the presence of a deeper normal fault. Hence, the thick-skinned tectonic influence mentioned by Cooper (2000) is confirmed in both the northern and southern seismic sections. However, the pop-up structure itself is a thin-skinned detachment structure (Thompson 1989) with a deeper influence, but the degree of normal fault re-activation is not as great as in Cooper's (2000) section, 35 km to the south. Normal faults in the Trutch-Halfway River area are cut and offset by the Besa River detachment (Figure 1-8). The Pink and Spruce Mountain structures owe their existence to a combination of Thompson's (1989) and Cooper's (2000) tectonic models.

REFERENCES

- Bally, A.W., Gordy, P.L., and Stewart, G.A., 1966. Structure, seismic data, and orogenic evolution of southern Canadian Rocky Mountains. *Bulletin of Canadian Petroleum Geology*, v. 14, no.3, p. 337-381.
- Bamber, E.W. and Mamet, B., 1978. Carboniferous biostratigraphy and correlation, north-eastern British Columbia and southwestern district of MacKenzie. *Geological Survey of Canada Bulletin* 266, 65 p.
- Bamber, E.W., Taylor, G.C. and Procter, R.M., 1968. Carboniferous and Permian stratigraphy of northeastern British Columbia. *Geological Survey of Canada Paper* 68-15, 25 p.
- Barclay, J.E. 1988. The lower Carboniferous Golata Formation of the Western Canada Basin, in the context of sequence stratigraphy. *Canadian Society of Petroleum Geologists Memoir* 15, p. 1-14.
- Barclay, J.E., Krause, F.F., Campbell, R.I., Utting, J., 1990. Dynamic casting and growth faulting. Dawson Creek Graben Complex, Carboniferous-Permian Peace River Embayment, Western Canada. *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 115-145.
- Barnes, C.R. and Pyle, L.J., 2000. Lower Devonian stratigraphic framework along a MacDonald Platform to Ospika Embayment transect, Trutch, Halfway River and Ware map areas (94G, B, F), northeastern British Columbia. In: *Central Foreland NATMAP Project 2000 Fall Workshop - Program and Abstracts*, MacNaughton R. (ed.), Geological Survey of Canada, 43 p.
- Bégin, N.J. and Spratt, D.A., 2002. Role of transverse faulting in along-strike termination of Limestone Mountain Culmination, Rocky Mountain thrust and fold belt, Alberta. *Journal of Structural Geology*, v. 24, p. 689-707.
- Belyea, H.R. and Norris, A.W., 1962. Middle Devonian and older Paleozoic formations of southern district of MacKenzie and adjacent areas. *Geological Survey of Canada Paper* 62-15, 82 p.
- Bloy, G. and Scott, B., 1993. Characterization of the Belloy Formation Reservoir, Boundary Lake Pool, northeastern British Columbia. In: *Carboniferous to Jurassic Pangea, first international symposium - Program and Abstracts*. Glass D. (ed.), McCara Printing Limited, 27 p.
- Boyer, S.E. and Elliott, D., 1982. Thrust systems. *Bulletin of the American Association of Petroleum Geologists*, v. 66, no. 9, p. 1196-1230.

Cant, D.J., 1988. Regional Structure and development of the Peace River Arch, Alberta. A Paleozoic failed-rift system? *Bulletin of Canadian Petroleum Geology*, v. 36, p. 284-295.

Cecile, M.P., 1997. Geology, Sikanni Chief River area, parts of NTS sheets 94G/3 (Marion Lake), 94G/6 (Mount Withrow). Geological Survey of Canada Open File 3471, scale 1:50,000.

Cecile, M.P., 1999. Preliminary Geology of the Marion Lake map area (94G/03). Geological Survey of Canada Open File 3736, scale 1:50,000.

Cecile, M.P. and Legun, A., 2001. Preliminary Geology of the Mount McCusker map area (94G/04), Geological Survey of Canada Open File 3767, scale 1:50,000.

Cecile, M.P., Morrow, D.W., Williams, G.K., 1997. Early Paleozoic (Cambrian to Early Devonian) tectonic framework, Canadian Cordillera. *Bulletin of Canadian Petroleum Geology*, v. 45. no. 1, p. 54-74.

Cecile, M.P. and Norford, B.S., 1979. Basin to Platform transition, lower Paleozoic strata of Ware and Trutch map areas, northeastern British Columbia. In: *Current Research, Part A Geological Survey of Canada Paper 79-1A*, p. 219-226.

Chamney, T.P., 1978. Albian Foraminifera of the Yukon Territory. Geological Survey of Canada, *Bulletin* 253, 62 p.

Cobbold, P.R., 1975. Fold propagation in single embedded layers. *Tectonophysics*, v. 27, p. 333-351.

Cobbold, P.R. and Gapais, D., 1987. Shear criteria in rocks: an introductory review. *Journal of Structural Geology*, v. 9, no. 5, p. 521-523.

Cook, F., 1993. Proterozoic crustal transition beneath the Western Canada Sedimentary basin. *Geology*, v. 21, p. 785-788.

Cooper, M. 2000. Structural style variations in the BC foothills. In: *GeoCanada 2000 – The Millennium summit, Programs and Abstracts*, Rod Blais (coord.), Geological Association of Canada, Mineralogical Association of Canada, Canadian Well Logging Society, Canadian Geophysical Union, Canadian Society of Exploration Geophysicists, Canadian Society of Petroleum Geologists. Iron Leaf Communications, CD-Rom.

Dahlstrom, C.D.A., 1970. Structural geology in the eastern margin of the Canadian Rocky Mountains. *Bulletin of Canadian Petroleum Geology*, v. 18, p. 332-406.

Davies, G.R., 1997. The upper Triassic Baldonnel and Pardonet formations, Western Canada Sedimentary Basin. *Bulletin of Canadian Petroleum Geology*, v. 45, no. 4, p. 643-674.

Davies, G.R., Moslow, T.F., and Sherwin, M.D., 1997. The Lower Triassic Montney Formation. *Bulletin of Canadian Petroleum Geology*, v. 45 no. 4, p. 474-505.

Dixon, J., 1999. Isopach maps of Triassic units in the Trutch (94G) map area (Central Forelands NATMAP). Geological Survey of Canada Open File 3765, 21p., 9 maps.

Evoy, R.W., 1997. Lowstand shorefaces in the middle Triassic Doig Formation. *Bulletin of Canadian Petroleum Geology*, v. 45, no. 4, p. 537-552.

Fermor, P.R. and Moffat, I.W., 1992. Tectonics and structure of the Western Canada foreland basin. In: *Foreland Basins and Thrust Belts*, R. Macqueen and D. Leckie (eds.). American Association of Petroleum Geologists Memoir 55, p. 81-105.

Fox, F.G., 1959. Structure and accumulation of hydrocarbons in southern Foothills, Alberta, Canada. *Bulletin of the American Association of Petroleum Geologists*, v. 43, no. 5, p. 992-1025.

Gibson, D.W., 1971. Triassic stratigraphy of the Sikanni Chief River – Pine Pass region, Rocky Mountain Foothills, northeastern British Columbia. Geological Survey of Canada Paper 70-31, 105p.

Gibson, D.W., 1992. Stratigraphy, sedimentology, coal geology and depositional environment of the Lower Cretaceous Gething Formation, northeastern British Columbia and west-central Alberta. *Geological Survey of Canada Bulletin* 43, 127 p.

Gibson, D.W., and Edwards, D.E., 1990. An Overview of Triassic stratigraphy and depositional environments in the Rocky Mountain Foothills and Western Interior Plains, Peace River Arch area, northeastern British Columbia. *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 146-158.

Glass, D., 1997. *Lexicon of Canadian Stratigraphy volume 4: Western Canada*, D. Glass (ed.). Canadian Society of Petroleum Geologists.

Guardia, F.J., 1971. Report on the coal deposits of Pink Mountain, British Columbia. British Columbia Department of Mines and Energy Open File Report 36, 23 p.

Hage, C.O., 1944. Geology adjacent to the Alaska Highway between Fort St. John and Fort Nelson, BC. Geological Survey of Canada Paper 44-30, 22 p.

Hunt, A.D. and Ratcliffe, J.D., 1959. Triassic stratigraphy, Peace River area, Alberta and

British Columbia, Canada. Bulletin of the American Association of Petroleum Geologists, v. 43, no. 3, p. 563-589.

Irish, E.J.W., 1963, Late Carboniferous and Permian stratigraphy of a part of northeastern British Columbia. Bulletin of Canadian Petroleum Geology, v.11, no. 4, p. 373-388.

Irish, E.J.W., 1970. Halfway River map-area, British Columbia. Geological Survey of Canada Paper 69-11.

Johns, M.J., Barnes, C.R., and Orchard, M.J., 2000. Progress Report on Middle to Late Triassic ichthyolith biostratigraphy and regional thermal maturation studies in the Trutch map area of northeastern British Columbia. In: Central Foreland NATMAP Project 2000 Fall Workshop - Program and Abstracts, MacNaughton R. (ed.), Geological Survey of Canada , 43 p.

Jones, P.B., 1971. Folded faults and sequence of thrusting in Alberta Foothills. Bulletin of the American Association of Petroleum Geologists, v. 55, no. 2, p. 292-306.

Kidd, F.A., 1963. The Besa River Formation. Bulletin of Canadian Petroleum Geology, v.11, no. 4, p. 369-372.

Lageson, D.R., 1984. Structural geology of Stewart Peak culmination, Idaho-Wyoming thrust belt. Bulletin of the American Association of Petroleum Geologists, v.68, no. 4, p. 401-416.

Lane, L.S., 2001. Geology of the Redfern Lake map area (94G/5), Geological Survey of Canada Open File 4172, scale 1:50 000, 1 sheet.

Lane, L.S., Cecile, M.P., Currie, L.D., and Stockmal, G.S., 1999. Summary of fieldwork in Trutch and Toad River map areas, Central Foreland NATMAP project, northeastern British Columbia. In: Current Research 1999-E, Geological Survey of Canada, p. 1-8.

Law, J., 1955. Geology of northwestern Alberta and adjacent areas. Bulletin of the American Association of Petroleum Geologists, v. 39, no. 10, p. 1927-1978.

Law, J., 1981. Mississippian correlations, northeastern British Columbia, and implications for oil and gas Exploration. Bulletin of Canadian Petroleum Geology, v. 29, no. 3, p. 378-398.

Leggett, S., Young, R. and Eibenschutz, D., 1993. Facies of the Belloy Formation (Lower Permian), northeastern British Columbia. In: Carboniferous to Jurassic Pangea, first international symposium; program and abstracts. Glass D. (ed.). McCara Printing Limited, 27 p.

- Lepard, D., Twelker, S., Tzeng, P., Mottershead, K., Palmer, B., Slater, J., and Lai, G., 1999. Sample and Analysis Management System (SAMS), version 3.10. Geological Survey of Canada.
- Liu, S. and Dixon, J.M., 1992. Centrifuge modelling of thrust faulting: structural variation along strike in fold-thrust belts. *Tectonophysics*, v. 188, p. 39-62.
- McLearn, F.H. and Kindle, E.D., 1950. Geology of northeastern British Columbia. Geological Survey of Canada Memoir 259, 236 p.
- McMechan, M.E., 1990. Upper Proterozoic to Middle Cambrian history of the Peace River Arch: evidence from the Rocky Mountains, *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 36-44.
- McNeil, D.H., 2000. Micropalaeontology report on forty-three outcrop samples from the upper Jurassic and Lower Cretaceous of the Pink Mountain area, Trutch corridor, northeastern British Columbia (NTS 94G). Geological Survey of Canada Internal Report 1-DHM-2000, 9 p.
- Nicholls, E., Brinkman, D. and Wu, X., 1998. A new Archosaur from the Upper Triassic Pardonet Formation of British Columbia. *Canadian Journal of Earth Sciences*, v. 35, p. 1134-1142.
- Norford, B.S., 1990. Ordovician and Silurian stratigraphy, paleogeography and depositional history in the Peace River Arch area, Alberta and British Columbia. *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 45-54.
- O'Connell, S.C., Dix, G.R. and Barclay, J.E., 1990. The origin, history, and regional structural development of the Peace River Arch, Western Canada. *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 4-24.
- Pelletier, B.R. 1964. Triassic stratigraphy of the Rocky Mountains between Peace and Muskwa rivers, northeastern British Columbia. Geological Survey of Canada Paper 63-33, 89 p.
- Pelletier, B.R. and Stott, D.B., 1963. Trutch map area, British Columbia 94G. Geological Survey of Canada Paper 63-10, 12 p.
- Poulton, T.P., Titterton, J., and Dolby, G., 1990. Jurassic strata of northwestern (and west-central) Alberta and northeastern British Columbia. *Bulletin of Canadian Petroleum Geology*, v. 38A, p. 159-175.

Province of British Columbia Energy, Mines and Petroleum Resources, 1979. Trutch-Beaton River – oil and gas fields, British Columbia Energy, Mines and Petroleum Resources Regional map A4, scale 1:200,000.

Pugh, D.C., 1975. Cambrian Stratigraphy from Western Alberta to northeastern British Columbia. Geological Survey of Canada Paper 74-37, 31 p.

Pyle, L.J. and Barnes, C.R., 2000. Upper Cambrian to Lower Silurian stratigraphic framework of platform-to-basin facies, northeastern British Columbia. *Bulletin of Canadian Petroleum Geology*, v. 48, no. 2, p. 123-149.

Ross, G.M., McNicoll, V.J., Geldsetzer, H.H.J., Parrish, R.R., Carr, S.D. and Kinsman, A., 1993. Detrital zircon geochronology of Siluro-Devonian sandstones, Rocky Mountains, northeastern British Columbia. *Bulletin of Canadian Petroleum Geology*, v. 41 n. 3, p. 349-357.

Schroder-Adams, C.J., 1999. Cretaceous stratigraphy and sea-level history in northeastern British Columbia. A high accommodation setting. In: Central Foreland NATMAP Project 1999 Fall Workshop, Programs and Abstracts, L. Lane (ed.). Geological Survey of Canada, 51 p.

Stockmal, G.S., 1999. Abrupt stratigraphic thickness variations and associated changes in structural style across cross-strike lineaments in the northeastern B.C. Foothills. evidence from bedrock mapping of pre-existing growth faults and implications for exploration. In: Central Foreland NATMAP Project 1999 Fall Workshop, Programs and Abstracts, L. Lane (ed.). Geological Survey of Canada, 51 p.

Storrs, G.W., 1991. Note on a second occurrence of *Thalattosaurus* remains (Reptilia. Neodiapsida) in British Columbia. *Canadian Journal of Earth Sciences*, v. 28, p. 2065-2068.

Stott, D.F., 1982. Lower Cretaceous Fort St. John Group and Upper Cretaceous Dunvegan Formation of the Foothills and Plains of Alberta, British Columbia, District of MacKenzie and Yukon Territory. Geological Survey of Canada, Bulletin 328.

Stott, D.F., 1991. Geotectonic correlation chart, Sheet 2, Prairie Provinces and British Columbia. In: *Sedimentary Cover of the North American Craton*. Canada, D.F. Stott and J.D. Aiken (eds.). Geological Survey of Canada, *Geology of Canada*, no. 5., 825 p.

Stott, D.F., 1998. Fernie Formation and Minnes Group (Jurassic and Lowermost Cretaceous), northern Rocky Mountain Foothills, Alberta and British Columbia. Geological Survey of Canada, Bulletin 516.

Stott, D.F., Goruk, G.L. and Pelletier, B.R., 1963. Trutch, British Columbia 94G, Geological Survey of Canada Preliminary Map 12-1963.

Tappan, H., 1955. Foraminifera from the Arctic Slope of Alaska. Part 2, Jurassic foraminifera. United States Geological Survey, Professional Paper 236-B, p. 21-90.

Taylor, G.C. 1979. Geology of east Ware (94F) and west Trutch (94G) map areas 1:250,000 Scale. Geological Survey of Canada Open File Report 606.

Taylor, G.C., MacIntyre, D.G., Okulitch, A.V., Cullen, B., Massey, N., and Bellefontaine, K., (compilers) 1999. Geology, Fort Nelson British Columbia, Central Foreland NATMAP NO-10-G, Scale 1:500,000, Geological Survey of Canada Open File 3604 (rev).

Taylor, G.C. and MacKenzie, W.S. 1970. Devonian Stratigraphy of northeastern British Columbia. Geological Survey of Canada, Bulletin 186.

Then, D.R. and Dougherty, B.J., 1983. A new procedure for extracting foraminifera from indurated organic shale. In: Current Research, Part B, Geological Survey of Canada, Paper 83-1B, p. 413-414.

Thompson, R.I., 1989. Stratigraphy, tectonic evolution and structural analysis of the Halfway River map-area (94 B), Northern Rocky Mountains, British Columbia. Geological Survey of Canada, Memoir 425.

Tozer, E.T., 1982. Late Triassic (Upper Norian) and earliest Jurassic (Hettangian) rocks and Ammonoid faunas, Halfway River and Pine Pass map areas, British Columbia. In: Current Research, Part A, Geological Survey of Canada, Paper 82-1A, p. 385-391.

Trotter, R., 1989. Sedimentology and depositional setting of the Granite Wash of the Utikuma and Red Earth areas, north-central Alberta. Unpublished MSc. Thesis, Dalhousie University, 378 p.

Wilson, R.C.L. and Williams, C.A., 1979. Oceanic transform structures and the development of Atlantic continental margin sedimentary basins – a review. Journal of the Geological Society of London, v. 136, p. 311-320.