

Homogeneous Coordinates

Jules Bloomenthal and Jon Rokne
Department of Computer Science
The University of Calgary

Introduction

Homogeneous coordinates have a natural application to Computer Graphics; they form a basis for the projective geometry used extensively to project a three-dimensional scene onto a two-dimensional image plane. They also unify the treatment of common graphical transformations. The graphical use of homogeneous coordinates is due to [Roberts, 1966]; today, homogeneous coordinates are presented in many computer graphics texts (such as [Foley, Newman, Rogers]); in particular, [Newman] provides an appendix of homogeneous techniques. [Riesenfeld] provides an excellent introduction to homogeneous coordinates and their algebraic, geometric and topologic importance to Computer Graphics. [Bez] further discusses their algebraic and topological properties and [Blinn77, Blinn78] develop additional applications for Computer Graphics.

Homogeneous coordinates are also used in the related areas of CAD/CAM [Zeid], robotics [McKerrow], surface modeling [Farin], and computational projective geometry [Kanatani]. They can also extend the number range for fixed point arithmetic [Rogers].

Our aim here is to provide an intuitive yet theoretically based discussion that assembles the key features of homogeneous coordinates and their applications to Computer Graphics. These applications include affine transformations, perspective projection, line intersections and clipping. For the sake of clarity in accompanying illustrations, we confine our development to two dimensions and then use the intuition gained to present the use of homogeneous coordinates in three dimensions.

[Kline] provides a brief history of homogeneous coordinates, crediting Möbius with their introduction. Given a fixed triangle in the plane, Möbius defined a set of homogeneous coordinates for a point p to be the weights required at the triangle vertices such that p becomes the center of gravity of the triangle (Figure 1, left). The point p is computed as

$$\mathbf{p} = (w_a \mathbf{a}, w_b \mathbf{b}, w_c \mathbf{c}),$$

with the condition that $w_a + w_b + w_c = 1$; this is an easily solved system of equations. The three unknowns w_a , w_b , and w_c , are called the *barycentric coordinates* of p with respect to \mathbf{a} , \mathbf{b} , and \mathbf{c} . [Farin] relates w to area by $w_a = A_b / (A_a + A_b + A_c)$ and similarly for w_b and w_c (Figure 1, middle).

Plücker defined another set of homogeneous coordinates by considering the signed distances from a point to the edges of a fixed triangle (Figure 1, right); here, $\mathbf{p} = (l_a, l_b, l_c)$.




Figure 1: barycentric and Plücker coordinates.

Barycentric and Plücker coordinates are both examples of coordinate systems in which $n+1$ values represent an n -dimensional point. One immediate attribute of these coordinate systems is the invariance of a point when scaled: scaling the Möbius weights or the size of the Plücker triangle does not change the position of p . For an exposition of n -dimensional Plücker coordinates (also known as Grassman coordinates), see [Stolfi].

Homogeneous Coordinates for the Projective Plane

The homogeneous coordinates commonly used in Computer Graphics are equivalent to Plücker coordinates in which one side of the triangle is at infinity and the opposite vertex is the origin of the coordinate system. The plane containing the triangle with the line at infinity is known as the *projective plane*.

[Aleksandrov] describes the projective plane by considering all lines and planes passing through a given point s ; if they are intersected by a plane P that does not pass through s , then each point (or line) on P may be associated with a line (or plane) through s , as shown in Figure 2, left. This does not quite imply a one-to-one mapping of lines (or planes) through s with points (or lines) on P , because those lines or planes through s parallel to P do not intersect P . By convention, however, the parallel lines are said to intersect P at *ideal* (infinitely distant) points; the parallel plane is said to intersect P at the *ideal line*. The plane P , augmented by ideal points and the ideal line, is known as the projective plane. It cannot be represented within the finite Euclidean coordinate system; it can be, however, be represented by homogeneous coordinates, and this is the fundamental reason for their use in projective geometry.




Figure 2: the projective plane.

Projective geometry is, in a sense, the geometry of imaging. This was already understood, for example, by Albrecht Dürer who used a mechanical device to draw objects in perspective [Penna]. It is, therefore, a natural tool for Computer Graphics. General projective geometry is discussed in several texts [Coxeter, Ryan], and its use in Computer Graphics is discussed in [Herman, Penna].

Mathematically, the mapping from planes and lines through s to lines and points on the projective plane is the transformation of the usual Euclidean space into projective space. The following three statements each define the two-dimensional projective space, ρ^2 (from [Ryan]):

- 1) The set of all equivalence classes of ordered triples of non-zero vectors in \mathcal{E}^3 , where equivalence is the mutual proportionality of two vectors.
- 2) The set of all lines passing through the origin of \mathcal{E}^3 .
- 3) The set of all pairs of antipodal points of S^2 , the unit sphere in \mathcal{E}^3 .

We denote the usual two-dimensional Euclidean space (also known as ‘physical’ or ‘proper’ space) by \mathcal{E}^2 . Within \mathcal{E}^2 , each point is represented as a two-component vector, (x', y') , where both x' and y' are finite values in an orthogonal coordinate system. The relationships between \mathcal{E}^3 , the unit sphere (\mathcal{S}^2), and the projective space (ρ^2), are illustrated in Figure 2. Points in ρ^2 are lines through the origin, as in the second definition above. This means that we use equivalence classes of coordinates of \mathcal{E}^3 to represent points in ρ^2 .

In practice, homogeneous coordinates represent ρ^2 by mapping each Euclidean point $(x', y') \in \mathcal{E}^2$ to $[x, y, w] \in \mathcal{E}^3$ ($w \neq 0$), which is a member of the equivalence class of points in ρ^2 . The mapping is achieved by the equivalences $x' \sim x/w$ and $y' \sim y/w$ (we enclose Euclidean coordinates within parentheses and homogeneous coordinates within brackets; their equivalence is signified by \sim).

ρ^2 is not a vector space in the same manner as \mathcal{E}^2 . Indeed, scalar multiplication simply forms a new representative of the equivalence class. Furthermore, if the representatives are chosen from the intersection with the plane $w = 1$, then the vector addition of two points in ρ^2 computes a representative of the midpoint between the two vectors.

Because of the division by w , the conversion of a homogeneous point to its Euclidean equivalent is inherently a projection of the homogeneous point onto the $w = 1$ plane. Figure 2, right, illustrates this projection in the two-dimensional case; three-dimensional homogeneous points on the \mathcal{S}^2 sphere are projected onto the $w = 1$ plane. [Riesensfeld] provides an illustration of four-dimensional homogeneous points projected onto a three-dimensional hyper-plane.

Homogeneous points represent projection and as a consequence can also represent points at infinity. Consider a homogeneous point as w approaches 0; for example, in Figure 3, $[2, 3, w]$ is shown for $w = (4, 2, 1, 1/2, 0)$. As w approaches 0, the projected Euclidean points move away from the origin in the $(2, 3)$ direction. At $w = 0$, the point is infinitely far and may be treated as a positionless vector.

Historically, the need for points at infinity arose from the work of Kepler and Desargues between the 16th to 17th centuries when they both realized that a parabola has two foci, one finite and one infinite [Coxeter].




Figure 3: projection to the $w = 1$ plane.

We close this section with a brief reflection on the term *homogeneous*, which [Oxford] defines as:

- 1) *Of the same kind so as to be commensurable.*
- 2) *Of the same degree or dimension: consisting of terms of the same dimension.*

With this in mind, consider a general ellipse in \mathcal{E}^2 :

$$f(x', y') = ax'^2 + by'^2 + cx'y' + dx' + ey' + f = 0.$$

This can be formulated homogeneously by replacing x' with x/w and y' with y/w ; multiplying by w^2 yields:

$$f'(x, y, w) = ax^2 + by^2 + cxy + dxw + eyw + fw^2 = 0.$$

Here, all terms have total variable degree two, as in definition 2), above. In fact, any polynomial function in \mathcal{E}^2 has a form equivalent to f and may be transformed, via a change to homogeneous coordinates, to a polynomial function with constant total degree for each term. This holds for any polynomial function in a finite dimensional space.

Homogeneous Coordinates for Two Dimensions

An important, practical aspect of the homogeneous coordinate system is its unification of the translation, scaling and rotation of geometric objects. In \mathcal{E}^2 Euclidean space, the most general affine mapping is

$$\mathbf{p}' = \mathbf{p}A + \mathbf{c}, \text{ or } (p'_x, p'_y) = (p_x, p_y)A + (c_x, c_y),$$

where \mathbf{p}' and \mathbf{p} are points in \mathcal{E}^2 , A is a 2×2 matrix representing scaling and rotation, and the point \mathbf{c} represents translation. In this formulation, translation is treated differently from rotation or scaling.

If, however, \mathbf{p} is represented as a homogeneous point $[p_x \ p_y \ 1]$ and a 2×3 matrix is employed, then translation may be treated the same as rotation and scaling:

$$[p'_x \ p'_y] = [p_x \ p_y \ 1] \begin{bmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \\ c_1 & c_2 \end{bmatrix}.$$

A more consistent, and ultimately simpler result is obtained by introducing a third column to the matrix and a third (homogeneous) coordinate to the result:

$$[p'_x \ p'_y \ 1] = [p_x \ p_y \ 1] \begin{bmatrix} a_{00} & a_{01} & 0 \\ a_{10} & a_{11} & 0 \\ c_1 & c_2 & 1 \end{bmatrix}.$$

$p'_x \ p'_y$ are unchanged, but the result is now three-dimensional and the matrix is square. Thus:

- the resulting (homogeneous) vector has the same dimension as the input (homogeneous) vector
- the matrix is invertible (assuming a non-zero determinant)
- two or more transformations may be concatenated.

The above transformation may be represented compactly by *bordering* the matrix A with the column vector $(0, 1)^T$ and the row vector \mathbf{c} :

$$[\mathbf{p}' \ 1] = [\mathbf{p} \ 1] \begin{bmatrix} A & 0 \\ \mathbf{c} & 1 \end{bmatrix}.$$

Here $[p_x' \ p_y' \ 1]$ and $[p_x \ p_y \ 1]$ are replaced with their compact forms $[p' \ 1]$ and $[p \ 1]$ so that the components conform in size with the sub-matrices. In this formulation, *all affine transformations are matrix multiplications*. Computer graphic transformations are usually encoded as matrices because of their representational simplicity and computational efficiency. We now summarize the use of homogeneous coordinates for affine transformations.

Transformations

- **Translation.** In the Euclidean coordinate system, translation is $p' = p + c = pI + c$, where I is the identity matrix. In the homogeneous coordinate system this is

$$[p_x' \ p_y' \ 1] = [p_x \ p_y \ 1] \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ c_x & c_y & 1 \end{bmatrix}.$$

Compact notation yields $[p' \ 1] = [p \ 1] \begin{bmatrix} I & 0 \\ c & 1 \end{bmatrix}$.

- **Rotation.** In the Euclidean coordinate system, rotation is defined by $p' = pR$,

$$\text{where } R = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}.$$

Objects are rotated by θ radians counter-clockwise about the origin. Using homogeneous coordinates, this becomes

$$[p_x' \ p_y' \ 1] = [p_x \ p_y \ 1] \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

and, compactly, $[p' \ 1] = [p \ 1] \begin{bmatrix} R & 0 \\ 0 & 1 \end{bmatrix}$.

- **Scaling.** A non-uniform scaling is defined by $S = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}$.

In a homogeneous coordinate system this becomes

$$[p_x' \ p_y' \ 1] = [p_x \ p_y \ 1] \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

and, compactly, $[p' \ 1] = [p \ 1] \begin{bmatrix} S & 0 \\ 0 & 1 \end{bmatrix}$.

Sequences

A sequence, or concatenation, of affine transformations can be expressed as a single matrix. For example, a translation followed by a rotation and then a scaling is equivalent to

$$[\mathbf{p}' \ 1] = [\mathbf{p} \ 1] \begin{bmatrix} I & 0 \\ \mathbf{c} & 1 \end{bmatrix} \begin{bmatrix} R & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} S & 0 \\ 0 & 1 \end{bmatrix} = [\mathbf{p} \ 1] \begin{bmatrix} RS & 0 \\ \mathbf{c}RS & 1 \end{bmatrix}.$$

Reducing a sequence of transformations to a single matrix improves computational performance, especially when numerous points are to be transformed by the same transformation.

Homogeneous Lines

A line l in \mathcal{E}^2 is defined by $ax+by+c=0$, with a, b, c constant; the intersection of lines l_1 and l_2 is found by solving

$$\begin{aligned} a_1x+b_1y+c_1 &= 0 \\ a_2x+b_2y+c_2 &= 0, \end{aligned}$$

obtaining: $x = \begin{vmatrix} -c_1 & b_1 \\ -c_2 & b_2 \end{vmatrix} \Big/ \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}; y = \begin{vmatrix} a_1 & -c_1 \\ a_2 & -c_2 \end{vmatrix} \Big/ \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix}.$

The two lines l_1 and l_2 intersect in \mathcal{E}^2 provided they are not parallel. This special provision may be eliminated by representing a line homogeneously [Newman]:

$$ax+by+cw = 0 \text{ (with } w = 1, \text{ in general),}$$

and representing the intersection as a homogeneous point:

$$[x, y, w] = \left(\begin{vmatrix} -c_1 & b_1 \\ -c_2 & b_2 \end{vmatrix}, \begin{vmatrix} a_1 & -c_1 \\ a_2 & -c_2 \end{vmatrix}, \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \right).$$

Dividing by w yields a finite point in \mathcal{E}^2 if $w \neq 0$, and a point at infinity if $w = 0$ (ie, the lines are parallel). $[x, y, w]$ is simply the cross product, $[a_1, b_1, c_1] \times [a_2, b_2, c_2]$.

If the intersection of two lines is the cross product of their coefficients, the intersection of two parallel lines may be given as $[a, b, c] \times [a, b, d] = [b, -a, 0]$, or $[1, -a/b, 0]$, which is the point at infinity in the direction of the line's slope. Note that the line at infinity may be represented as $[0, 0, a]$; the dot product with this line and any point at infinity is zero; thus, all points at infinity lie on the line at infinity.

We also note the duality between line and point in that the cross product of two homogeneous points yields the coordinates of their connecting line. This duality between point and line in two dimensions has, in three dimensions, a corresponding duality between point and plane. See [Blinn77] for the use of homogeneous coordinates to represent lines in \mathcal{E}^3 .

Applications in Three Dimensions

We now apply homogeneous coordinates to three-dimensional Euclidean points. Early in the development of Computer Graphics, L. G. Roberts noted the value of homogeneous coordinates, stating, “the use of homogeneous coordinates throughout is extremely important in order to maintain the simplicity of the results, although its original purpose was to allow perspective transformations” [Roberts].

In other words, the use of the additional, homogeneous coordinate not only produces polynomials of fixed degree, as observed in the Introduction, it also provides a method for consistent manipulation of the Euclidean space.

It is well known that a 3×3 matrix can represent three-dimensional scaling and rotation, but not translation. As in the two-dimensional case, translation becomes possible with the addition of a row and column to the matrix:

$$[x' y' z' w'] = [x y z 1] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ t_x & t_y & t_z & 1 \end{bmatrix}$$

Although it is possible to represent this in compact form:

$$[p' 1] = [p 1] \begin{bmatrix} I & 0 \\ t & 1 \end{bmatrix}, \text{ where } I \text{ is the identity matrix and } t \text{ the translation,}$$

we do not do so when considering three-dimensional applications because individual components of the transformation matrix, will be discussed in detail.

Perspective Projection

Fundamental to three-dimensional Computer Graphics is the projection of three-dimensional objects onto a two-dimensional image plane. This projection is usually a perspective projection with the center of projection being the point of view (eyepoint) and the central projector perpendicularly intersecting the projective (image) plane. [Carlbom] provides a review of planar projections. In creating shaded images, a perspective projection is not strictly necessary; ray tracing, for example, computes pixel values directly without a perspective transformation of the object. For line drawings and polygon rendering, however, the perspective transformation is essential.

As shown in Figure 4, we identify the screen of an output device with a rectangular domain in \mathcal{E}^2 ; the rendering of objects in \mathcal{E}^3 is naturally performed via their projection to \mathcal{E}^2 .




Figure 4: perspective projection onto the raster-screen.

Referring to the side view in Figure 5, the image point (x', y') is computed from the object point (x, y, z) via similar triangles:

$$y'/y = D/z \text{ and } x'/x = D/z, \text{ or } (x', y') = (xD/z, yD/z).$$




Figure 5: the perspective projection

The division by z is known as the *perspective divide*. This non-linear relationship in three dimensions can be formulated as a linear relationship in four dimensions through the use of homogeneous coordinates; the formal foundation for such a formulation is provided by [Bez].

Until now we have used the homogeneous coordinate to accommodate the larger matrix required to perform affine transformations; after the transformation the fourth coordinate, w , has remained 1. If, however, the resulting homogeneous coordinate w is proportional to the distance from the eye to a point, a perspective projection of that point onto the image plane is effected.

For example, if we assume the eye to be located at the origin and directed towards the positive z -axis, with the x -axis to the right and the y -axis up, we can project points onto the $z = D$ plane with the following matrix transformation:

$$(x', y', z') = (xD/z, yD/z, D) \sim [x \ y \ z \ z/D] = [x \ y \ z \ 1] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1/D \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

As D approaches ∞ , $w' = 1/D$ approaches 0 and the transformed points become infinitely far; their position parallel to the xy plane is unaffected by z and, effectively, an orthographic transformation results.

Perspective, Projection, and Perspective Projection

The perspective projection above produces a constant $z' = D$; this loss of depth information is due to the linear dependence of the matrix's third and fourth columns. If we introduce a second non-zero term into the third column, eg. -1 , we do not affect x' and y' , but z' becomes $D-D/z$. The motivation for this additional term is to compress the Euclidean space $z \in [1, \infty]$ to $z' \in [0, D]$.

$$(x', y', z') = (xD/z, yD/z, D-D/z) \sim [x \ y \ z^{-1} \ z/D] = [x \ y \ z \ 1] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1/D \\ 0 & 0 & -1 & 0 \end{bmatrix}.$$

This, then, is the perspective transformation; multiplied by a pure projection it yields the perspective-projection transformation:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1/D \\ 0 & 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & D & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1/D \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

perspective projection perspective-projection

Perspective Space

The homogeneous perspective transformation transforms Euclidean points in \mathcal{E}^3 (represented as homogeneous points) to new homogeneous points, which may then be converted to Euclidean space. These transformed points exist in *perspective space*. We now investigate the properties of the perspective transformation in terms of the relation between perspective space and the untransformed *object space*.

In an affine transform, the last column is $[0 \ 0 \ 0 \ 1]^T$; multiplication with a direction vector $[a, b, c, 0]$ yields another direction vector $[a', b', c', 0]$. In a perspective transform, however, the last column is $[0 \ 0 \ -1/D \ 1]^T$; multiplication with a direction vector yields $[a', b', c', -c/D]$. This is a homogeneous *point* not *direction* and is known as the *vanishing point* for the given direction. A set of three-dimensional parallel lines will intersect at the same vanishing point, with the one exception that parallel lines in the xy plane remain parallel after perspective.

In many graphics systems, points in object space are transformed by a matrix that is the concatenation of all rotation, scaling, translation, and perspective transformations. Usually the projection transformation is not incorporated into this concatenation, for several reasons. First, the perspective matrix is invertible whereas the perspective-projection matrix is singular. Secondly, the value of z' before projection is often useful for incremental scanline depth sorting or for intensity variation in line drawings. More importantly, however, z' can simplify three-dimensional clipping of Euclidean lines and line segments against the *viewing frustum* by clipping homogeneously against a parallelepiped. After clipping, points are simply projected to the $z = D$ image plane and displayed.

The viewing frustum consists of six planes defined in terms of the eye position, the visible portion of the image plane, and the allowable depths of three-dimensional objects. The viewing frustum shown in Figure 6 has a visible image sized $2*S_x$ by $2*S_y$, and the near and far clipping planes arbitrarily set to $z = D$ and $z = F$.




Figure 6: viewing frustum

To effect the image size above, we simply scale the first and second columns of the perspective matrix by S_x and S_y respectively. Assuming a square window, S_x and S_y may be derived from the field of view, fov , by $S_x = S_y = D \cdot \tan(fov/2)$.

To simplify clipping, we'd like the near clipping plane to transform to $z' = 0$, and the far clipping plane to transform to $z' = 1$. By considering the transformations of the points $(0, 0, D)$ and $(0, 0, F)$, where $[P]$ is the perspective matrix:

$$\begin{aligned} [0 \ 0 \ D \ 1][P] &= [0, 0, D-1, 1] \sim (0, 0, D-1) \\ [0 \ 0 \ F \ 1][P] &= [0, 0, F-1, F/D] \sim (0, 0, D-D/F) \end{aligned}$$

we see that the near and far clipping planes can be transformed by scaling the z-terms, resulting in:

$$[P] = \begin{bmatrix} 1/S_x & 0 & 0 & 0 \\ 0 & 1/S_y & 0 & 0 \\ 0 & 0 & \frac{1}{D(1-D/F)} & 1/D \\ 0 & 0 & \frac{-1}{1-D/F} & 0 \end{bmatrix}$$

Now, as desired:

$$\begin{aligned} [0 \ 0 \ D \ 1][P] &= \left[0, 0, \frac{D}{D(1-D/F)} - \frac{1}{1-D/F}, 1\right] = [0, 0, 0, 1] \\ [0 \ 0 \ F \ 1][P] &= \left[0, 0, \frac{F}{D(1-D/F)} - \frac{1}{1-D/F}, \frac{F}{D}\right] = \left[0, 0, \frac{F-D}{F(1-D/F)}, 1\right] = [0, 0, 1, 1]. \end{aligned}$$

Example Perspective Transformations

To familiarize ourselves with the perspective matrix, we list the following homogeneous points and their transformations:

$[a \ b \ D \ 1][\mathbf{P}] = [a', b', 0, 1]$	point on near clipping plane ($z = D$) transforms to xy plane ($z' = 0$)
$[a \ b \ F \ 1][\mathbf{P}] = [a', b', 1, 1]$	far plane transforms to $z' = 1$
$[0 \ 0 \ 1 \ 0][\mathbf{P}] = [0, 0, 1/(1-D/F), 1]$	vanishing point for line parallel to z -axis
$[a \ b \ c \ 0][\mathbf{P}] = [a', b', c', 1]$	vanishing point for arbitrary line
$[a \ b \ D \ 1][\mathbf{P}] = [a/S_x, b/S_y, 0, 1]$	but for scale, point on image plane unchanged
$[a \ b \ 0 \ 0][\mathbf{P}] = [a/S_x, b/S_y, 0, 0]$	point at infinity in xy plane is unchanged; <i>ie</i> , parallel lines in xy plane remain parallel
$[a \ 0 \ b \ 0][\mathbf{P}] = [\frac{aD}{bS_x}, 0, \frac{1}{1-D/F}, 1]$	point at ∞ in xz plane lies on perspective horizon
$[\pm S_x \ \pm S_y \ D \ 1][\mathbf{P}] = [\pm 1, \pm 1, 0, 1]$	frustum corners become image window corners
$[0 \ 0 \ 0 \ 1][\mathbf{P}] = [0, 0, \frac{-1}{1-D/F}, 0]$	eye transforms to point at infinity on negative perspective z -axis

Associated Vertex Transformations

The efficient shading of a polygon whose vertices have been transformed to the display screen is usually accomplished by incremental techniques described in the literature [Newman]. Similar techniques may be applied to vertex parameters such as color and texture. Any parameter interpolated across a polygon must follow the same transformation as the polygon vertices; if the transformation includes perspective, a homogeneous division is required at each pixel within the screen space polygon (in practice, linear interpolation in screen space is usually not objectionable for parameters such as color, but is immediately apparent when applied to texture coordinates).

Specifically, for perspective transformations, an interpolated parameter will have the form

$$(as_x + bs_y + c) / (ds_x + es_y + f),$$

where s_x and s_y are screen space coordinates. [Heckbert] refers to this as 'rational linear interpolation,' in a detailed discussion of texture coordinate transformations. [Blinn92], which provides an intuitive review of graphics transformations and homogeneous coordinates, refers to this as 'hyperbolic interpolation.' In the case where the polygon is parallel to the projection plane, $d = e = 0$ and the above equation reduces to linear interpolation.

Homogeneous Clipping

Three-dimensional line segments and polygon edges must be clipped to the viewing frustum. This should be performed in perspective space, after vertex colors have been computed; otherwise, clipping, by modifying a vertex's location, would modify its color. By clipping in perspective space the integrity of the polygon is maintained longer. Also, the equations for the clipping planes in homogeneous screen space are of the form $x = w$, whereas they are more complex in object space. [Blinn78 and Blinn91] develop this material in greater detail.

We use '**' to represent homogeneous coordinates before the conversion to Euclidean space. That is, $[x^*, y^*, z^*, w^*] = [x \ y \ z \ 1][\mathbf{P}]$ and $(x', y', z') = (x^*/w^*, y^*/w^*, z^*/w^*)$.

As shown in Figure 6, the viewing frustum is transformed by the perspective matrix to a parallelepiped in perspective space, and (x', y', z') is clipped such that $-1 \leq x', y' \leq 1$ and $0 \leq z' \leq 1$. Note that negative z values result in negative w values, with $z' = z^*/w^* > 0$. Thus, points behind the eye must be clipped before the homogeneous division. In summary, we test that:

for positive w :		for negative w :
$-w^* \leq x^*, y^*$	-- left, bottom --	$-w^* \geq x^*, y^*$
$x^*, y^* \leq w^*$	-- right, top --	$x^*, y^* \geq w^*$
$0 \leq z^*$	-- near --	$0 \geq z^*$
$z^* \leq w^*$	-- far --	$z^* \geq w^*$

Let us represent the line segment to be clipped as, $\mathbf{p}^* = \mathbf{p}_1^* + \alpha(\mathbf{p}_2^* - \mathbf{p}_1^*)$, $\alpha \in [0, 1]$, where \mathbf{p}_1^* and \mathbf{p}_2^* are the perspective-space endpoints $[x_1^*, y_1^*, z_1^*, w_1^*]$ and $[x_2^*, y_2^*, z_2^*, w_2^*]$. The intersection of the line segment with a clipping plane is then given in terms of α :

$$\begin{aligned} \alpha_{\text{left}} &= \frac{x_1^* + w_1^*}{-(w_2^* - w_1^*) - (x_2^* - x_1^*)} & \alpha_{\text{right}} &= \frac{x_1^* - w_1^*}{(w_2^* - w_1^*) - (x_2^* - x_1^*)} \\ \alpha_{\text{bottom}} &= \frac{y_1^* + w_1^*}{-(w_2^* - w_1^*) - (y_2^* - y_1^*)} & \alpha_{\text{top}} &= \frac{y_1^* - w_1^*}{(w_2^* - w_1^*) - (y_2^* - y_1^*)} \\ \alpha_{\text{near}} &= \frac{z_1^*}{-(z_2^* - z_1^*)} & \alpha_{\text{far}} &= \frac{z_1^* - w_1^*}{(w_2^* - w_1^*) - (z_2^* - z_1^*)} \end{aligned}$$

For example, consider the perspective space line segment from $\mathbf{p}_1^* = [2, y_1, z_1, 2]$ to $\mathbf{p}_2^* = [-1, y_2, z_2, 1/2]$. For the first point, $\mathbf{p}_{1x}^* > -\mathbf{p}_{1w}^*$ ($2 > -2$); but, for the second point, $\mathbf{p}_{2x}^* < -\mathbf{p}_{2w}^*$ ($-1 < -1/2$), implying an intersection with the left clipping plane. We compute

$$\alpha_{\text{left}} = (2+2)/((2-1/2)-(-1-2)) = 8/3.$$

The intersection with the left clipping plane is now given by

$$\mathbf{p}^* = [-2/3, y_1^* + (8/3)(y_2^* - y_1^*), z_1^* + (8/3)(z_2^* - z_1^*), 2/3].$$

The projected x -coordinate of $\mathbf{p}^* = x^*/w^* = (-2/3)/(2/3) = -1$, which is the left boundary of the viewing frustum.

Conclusion

In this paper we have offered a unified view of homogeneous coordinates within a Computer Graphics context. First, a brief historical review revealed that, as the understanding of perspective and projections increased, new coordinate systems were developed to represent the underlying spaces; one of these systems was the homogeneous coordinate system, which was later seen to possess properties useful for Computer Graphics.

Next, we formally introduced the homogeneous coordinate system. Its application in two-dimensional Euclidean space was discussed in some detail; it was shown that affine transformations can be effected consistently with matrix multiplication, thus simplifying sequences of transformations, and that the intersection of two-dimensional lines can be performed without special cases.

Homogeneous coordinates in three dimensions were discussed in greater detail, with particular attention devoted to perspective transformations. Finally, a method to clip lines with respect to the viewing frustum was provided.

Acknowledgements

We gratefully acknowledge Jim Blinn, Charlie Gunn, Paul Heckbert, and Ken Shoemake for numerous suggestions that improved this paper.

References

- Aleksandrov, A.D., Kolmogorov, A.N., Lavientèv, M.A. *Mathematics, its Content, Methods and Meaning*. MIT Press, 1969.
- Bez, H.E. *Homogeneous Coordinates for Computer Graphics*. Computer-Aided Design, 1983.
- Blinn, J. *A Homogeneous Formulation for Lines in 3 Space*. Proceedings of SIGGRAPH 77. In *Computer Graphics* 11 (3), 1977.
- Blinn, J. *A Trip Down the Graphics Pipeline: Line Clipping*. IEEE *Computer Graphics and Applications* 11 (1), 1991.
- Blinn, J. *Hyberbolic Interpolation*. IEEE *Computer Graphics and Applications* 12 (4), 1992.
- Blinn, J., Newell, M.E. *Clipping Using Homogeneous Coordinates*. Proceedings of SIGGRAPH 78. In *Computer Graphics* 12 (3), 1978.
- Carlbon, I. and Paciorek, J. *Geometric Projections and Viewing Transformations*. ACM *Computing Surveys* 10 (4), 1978.
- Coxeter, H.S.M. *Projective Geometry* (second edition), Springer-Verlag, New York, 1987.
- Farin, G. *Curves and Surfaces for Computer Aided Geometric Design* (second edition), Academic Press, New York, 1990.
- Foley, J.D., van Dam, A., Feiner, S.K., Hughes, J.F. *Computer Graphics, Principles and Practice* (second edition). Addison Wesley, Reading, Massachusetts, 1990.
- Heckbert, P. *Fundamentals of Texture Mapping and Image Warping*. Masters thesis, Computer Science Division, The University of California at Berkeley, May, 1989.
- Herman, I. *The Use of Projective Geometry in Computer Graphics*. Lecture Notes in Computer Science 564, Springer Verlag, 1991.
- Kantani, K. *Computational Projective Geometry*. *Computer Vision, Graphics, and Image Processing* 54, 1991.
- Kline, M. *Mathematical Thought from Ancient to Modern Times*. Oxford University Press, New York, 1972.
- Möbius, F. *Gesammelte Werke*, Volume 1, pp. 36-49, *Die Barycentrische Calcul*. Dr. M. Saendig oHG, Wiesbaden, Germany, 1967.
- Newman, W.M. and Sproull, R.F. *Principles of Interactive Computer Graphics*. McGraw-Hill, New York, 1979.
- Oxford University Press. *The Compact Edition of the Oxford English Dictionary*. Oxford University Press, New York, 1971.
- Penna, M.A., Patterson, R.R. *Projective Geometry and its Application to Computer Graphics*. Prentice Hall, Englewood Cliffs, N.J., 1986.

Riesenfeld, R.F. *Homogeneous Coordinates and Projective Planes in Computer Graphics*. IEEE Computer Graphics and Applications 1, 1981.

Roberts, L.G. *Homogeneous matrix representation and manipulation of n-dimensional constructs*. In: The Computer Display Review. C. W. Adams Associates, Inc., Cambridge, Massachusetts, 1966.

Rogers, D.F., Adams, J.A. *Mathematical Elements for Computer Graphics* (second edition). McGraw Hill, New York, 1990.

Ryan, P.J. *Euclidean and Non-Euclidean Geometry*. Cambridge University Press, Cambridge, 1986.

Stolfi, J. *Oriented Projective Geometry*. Academic Press, 1991.